

Table 3B
2007 Pregnancy Nutrition Surveillance ⁽¹⁾
North Carolina
Comparison of Racial and Ethnic Distribution by County

Comparison of Racial and Ethnic Distribution ⁽²⁾								
County	Number Included in Analysis	White, Not Hispanic %	Black, Not Hispanic %	Hispanic %	American Indian/Alaskan Native %	Asian/Pacific Islander %	Multiple Races %	All Other/Unknown %
Alamance	1,172	38.7	26.2	32.5	0.3	1.9	0.0	0.3
Alexander	246	82.5	5.7	8.5	0.4	2.4	0.0	0.4
Alleghany	54	*	*	*	*	*	*	*
Anson	232	25.0	69.8	3.4	0.4	1.3	0.0	0.0
Ashe	170	93.5	0.6	5.3	0.0	0.6	0.0	0.0
Avery	123	89.4	0.0	8.9	0.0	1.6	0.0	0.0
Beaufort	386	35.0	45.6	18.9	0.3	0.3	0.0	0.0
Bertie	175	14.3	84.0	1.1	0.6	0.0	0.0	0.0
Bladen	312	35.3	41.7	20.5	2.6	0.0	0.0	0.0
Brunswick	620	63.7	18.2	15.2	1.6	1.0	0.0	0.3
Buncombe	1,450	67.7	13.2	17.1	0.2	1.8	0.0	0.0
Burke	614	67.1	5.9	18.1	0.0	9.0	0.0	0.0
Cabarrus	1,233	40.0	22.1	36.8	0.0	1.1	0.0	0.0
Caldwell	548	81.2	7.1	10.8	0.4	0.2	0.0	0.4
Camden	31	*	*	*	*	*	*	*
Carteret	343	75.2	11.1	11.4	1.5	0.9	0.0	0.0
Caswell	119	49.6	40.3	10.1	0.0	0.0	0.0	0.0
Catawba	1,057	54.7	14.3	23.2	0.6	7.1	0.0	0.2
Chatham	350	32.9	15.7	50.6	0.0	0.3	0.0	0.6
Cherokee	139	88.5	2.9	4.3	2.9	0.7	0.0	0.7

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 3B
2007 Pregnancy Nutrition Surveillance ⁽¹⁾
North Carolina
Comparison of Racial and Ethnic Distribution by County

Comparison of Racial and Ethnic Distribution ⁽²⁾								
County	Number Included in Analysis	White, Not Hispanic %	Black, Not Hispanic %	Hispanic %	American Indian/Alaskan Native %	Asian/Pacific Islander %	Multiple Races %	All Other/Unknown %
Chowan	117	35.0	59.0	6.0	0.0	0.0	0.0	0.0
Clay	31	*	*	*	*	*	*	*
Cleveland	762	56.3	37.9	4.3	0.4	0.9	0.0	0.1
Columbus	436	40.4	45.4	11.5	2.8	0.0	0.0	0.0
Craven	912	53.1	33.4	9.5	0.4	3.5	0.0	0.0
Cumberland	3,356	38.9	44.1	12.2	2.2	2.5	0.0	0.1
Currituck	84	*	*	*	*	*	*	*
Dare	180	68.3	5.0	25.6	0.0	1.1	0.0	0.0
Davidson	1,037	64.4	14.5	17.9	0.7	2.3	0.0	0.2
Davie	209	70.8	7.2	22.0	0.0	0.0	0.0	0.0
Duplin	594	27.8	27.3	44.3	0.3	0.3	0.0	0.0
Durham	1,884	8.7	49.8	39.5	0.6	1.3	0.0	0.2
Edgecombe	620	19.7	74.8	5.5	0.0	0.0	0.0	0.0
Forsyth	2,431	23.2	37.9	38.0	0.2	0.7	0.0	0.0
Franklin	316	42.1	35.8	21.5	0.6	0.0	0.0	0.0
Gaston	1,520	56.4	27.1	14.9	0.1	1.1	0.0	0.4
Gates	30	*	*	*	*	*	*	*
Graham	75	*	*	*	*	*	*	*
Granville	325	34.8	44.9	18.8	0.0	1.5	0.0	0.0
Greene	221	25.3	50.7	23.5	0.5	0.0	0.0	0.0

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 3B
2007 Pregnancy Nutrition Surveillance ⁽¹⁾
North Carolina
Comparison of Racial and Ethnic Distribution by County

Comparison of Racial and Ethnic Distribution ⁽²⁾								
County	Number Included in Analysis	White, Not Hispanic	Black, Not Hispanic	Hispanic	American Indian/Alaskan Native	Asian/Pacific Islander	Multiple Races	All Other/Unknown
		%	%	%	%	%	%	%
Guilford	3,004	21.5	53.8	18.7	0.8	5.1	0.0	0.1
Halifax	551	24.0	67.9	2.7	5.3	0.2	0.0	0.0
Harnett	813	47.4	28.4	21.9	1.5	0.9	0.0	0.0
Haywood	318	85.2	2.8	10.1	1.6	0.3	0.0	0.0
Henderson	484	50.8	4.8	43.2	0.0	1.2	0.0	0.0
Hertford	204	15.7	81.9	2.0	0.5	0.0	0.0	0.0
Hoke	415	25.5	35.2	24.3	13.3	1.7	0.0	0.0
Hyde	30	*	*	*	*	*	*	*
Iredell	983	54.8	24.9	17.7	0.2	2.2	0.0	0.1
Jackson	255	60.4	2.4	11.0	23.5	2.7	0.0	0.0
Johnston	1,011	40.1	22.1	36.8	0.6	0.5	0.0	0.0
Jones	76	*	*	*	*	*	*	*
Lee	514	30.0	28.2	40.5	0.4	0.8	0.0	0.2
Lenoir	518	29.5	53.5	15.8	0.2	0.8	0.0	0.2
Lincoln	417	78.2	8.2	12.2	0.2	1.2	0.0	0.0
Macon	330	81.8	4.2	12.1	0.3	1.2	0.0	0.3
Madison	244	70.9	1.2	25.4	1.6	0.8	0.0	0.0
Martin	117	91.5	1.7	5.1	0.9	0.9	0.0	0.0
Mcdowell	195	31.8	60.5	6.7	0.0	1.0	0.0	0.0
Mecklenbur	4,647	13.0	52.3	30.9	0.4	3.1	0.0	0.2

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 3B
2007 Pregnancy Nutrition Surveillance ⁽¹⁾
North Carolina
Comparison of Racial and Ethnic Distribution by County

Comparison of Racial and Ethnic Distribution ⁽²⁾								
County	Number Included in Analysis	White, Not Hispanic	Black, Not Hispanic	Hispanic	American Indian/Alaskan Native	Asian/Pacific Islander	Multiple Races	All Other/Unknown
		%	%	%	%	%	%	%
Mitchell	88	*	*	*	*	*	*	*
Montgomery	239	41.8	27.2	28.5	0.4	2.1	0.0	0.0
Moore	292	49.0	28.8	19.9	1.4	1.0	0.0	0.0
Nash	736	29.1	56.3	13.6	0.7	0.3	0.0	0.1
New Hanove	1,166	41.6	36.0	21.1	0.3	0.6	0.0	0.4
Northhampt	163	20.9	76.1	2.5	0.0	0.6	0.0	0.0
Onslow	2,134	65.5	18.7	11.9	0.8	3.0	0.0	0.2
Orange	490	28.0	29.6	38.6	0.8	2.9	0.0	0.2
Pamlico	73	*	*	*	*	*	*	*
Pasquotank	280	33.6	51.8	13.6	0.0	1.1	0.0	0.0
Pender	351	51.3	25.4	22.8	0.3	0.3	0.0	0.0
Perquimans	68	*	*	*	*	*	*	*
Person	272	50.0	39.0	10.7	0.0	0.4	0.0	0.0
Pitt	1,226	29.0	59.2	10.7	0.2	0.8	0.0	0.1
Polk	54	*	*	*	*	*	*	*
Randolph	998	60.1	8.5	29.1	0.3	1.7	0.0	0.3
Richmond	528	47.0	36.2	13.8	2.3	0.8	0.0	0.0
Robeson	1,498	16.0	27.1	15.0	41.6	0.2	0.0	0.2
Rockingham	673	58.4	25.0	15.5	0.6	0.6	0.0	0.0
Rowan	977	48.8	26.4	23.4	0.1	1.0	0.0	0.2

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 3B
2007 Pregnancy Nutrition Surveillance ⁽¹⁾
North Carolina
Comparison of Racial and Ethnic Distribution by County

Comparison of Racial and Ethnic Distribution ⁽²⁾								
County	Number Included in Analysis	White, Not Hispanic %	Black, Not Hispanic %	Hispanic %	American Indian/Alaskan Native %	Asian/Pacific Islander %	Multiple Races %	All Other/Unknown %
Rutherford	395	70.4	17.7	11.1	0.0	0.8	0.0	0.0
Sampson	595	28.4	28.7	39.8	2.7	0.2	0.0	0.2
Scotland	443	29.1	54.2	0.7	15.6	0.2	0.0	0.2
Stanly	412	63.6	17.2	12.9	0.5	5.8	0.0	0.0
Stokes	252	90.9	4.0	4.8	0.0	0.4	0.0	0.0
Surry	509	68.2	3.7	27.1	0.4	0.6	0.0	0.0
Swain	155	58.7	0.6	1.9	36.1	2.6	0.0	0.0
Transylvan	169	79.9	8.3	10.1	0.0	1.2	0.0	0.6
Tyrrell	46	*	*	*	*	*	*	*
Union	1,014	37.2	24.6	37.7	0.0	0.5	0.0	0.1
Vance	496	27.4	57.3	13.9	0.4	1.0	0.0	0.0
Wake	4,418	19.4	39.6	38.3	0.2	2.2	0.0	0.2
Warren	167	14.4	72.5	9.6	3.0	0.6	0.0	0.0
Washington	147	25.2	68.7	6.1	0.0	0.0	0.0	0.0
Watauga	167	83.8	3.0	12.0	0.0	1.2	0.0	0.0
Wayne Coun	1,088	32.6	40.6	24.9	0.3	1.1	0.0	0.5
Wilkes	505	76.2	7.1	15.8	0.2	0.6	0.0	0.0
Wilson Cou	737	20.5	55.5	23.3	0.0	0.5	0.0	0.1
Yadkin	264	58.0	4.9	36.7	0.0	0.4	0.0	0.0
Yancey	104	89.4	0.0	10.6	0.0	0.0	0.0	0.0

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 3B
2007 Pregnancy Nutrition Surveillance ⁽¹⁾
North Carolina
Comparison of Racial and Ethnic Distribution by County

Comparison of Racial and Ethnic Distribution ⁽²⁾								
County	Number Included in Analysis	White, Not Hispanic %	Black, Not Hispanic %	Hispanic %	American Indian/Alaskan Native %	Asian/Pacific Islander %	Multiple Races %	All Other/Unknown %
Unknown Counties	259	41.7	33.6	20.8	3.1	0.8	0.0	0.0
North Carolina	64,288	39.9	33.9	22.4	2.0	1.8	0.0	0.1
Nation (Prior Year)	1,153,475	41.5	22.7	28.7	1.1	3.1	0.9	2.0

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

Table 4B
2007 Pregnancy Nutrition Surveillance ⁽¹⁾
North Carolina
Comparison of Age Distribution by County

Comparison of Age Distribution ⁽²⁾								
County	Number Included in Analysis	< 15 Years %	15 - 17 Years %	18 - 19 Years %	20 - 29 Years %	30 - 39 Years %	>= 40 Years %	Unknown %
Alamance	1,172	1.0	7.7	14.4	59.9	15.5	1.5	0.0
Alexander	246	0.0	8.1	17.9	58.5	13.8	1.6	0.0
Alleghany	54	*	*	*	*	*	*	*
Anson	232	0.9	12.9	17.2	53.9	15.1	0.0	0.0
Ashe	170	0.0	9.4	18.8	52.4	17.1	2.4	0.0
Avery	123	0.0	4.1	10.6	59.3	25.2	0.8	0.0
Beaufort	386	0.5	8.0	14.2	62.7	14.2	0.3	0.0
Bertie	175	0.6	8.6	20.6	59.4	10.9	0.0	0.0
Bladen	312	0.6	10.9	12.2	60.9	14.7	0.6	0.0
Brunswick	620	0.6	6.8	15.0	62.6	14.0	1.0	0.0
Buncombe	1,450	0.3	6.3	15.8	58.8	17.6	1.2	0.0
Burke	614	1.0	10.4	16.3	56.5	15.1	0.7	0.0
Cabarrus	1,233	0.4	7.5	14.8	58.0	18.4	0.9	0.0
Caldwell	548	0.0	10.8	19.0	56.4	13.0	0.9	0.0
Camden	31	*	*	*	*	*	*	*
Carteret	343	0.0	7.9	14.0	61.8	15.5	0.9	0.0
Caswell	119	0.0	8.4	8.4	67.2	16.0	0.0	0.0
Catawba	1,057	0.6	8.5	16.3	58.9	14.5	1.2	0.0
Chatham	350	0.3	10.3	12.0	58.6	16.9	2.0	0.0

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

Table 4B
2007 Pregnancy Nutrition Surveillance ⁽¹⁾
North Carolina
Comparison of Age Distribution by County

Comparison of Age Distribution ⁽²⁾								
County	Number Included in Analysis	< 15 Years %	15 - 17 Years %	18 - 19 Years %	20 - 29 Years %	30 - 39 Years %	>= 40 Years %	Unknown %
Cherokee	139	0.0	5.0	12.9	60.4	18.7	2.9	0.0
Chowan	117	0.9	9.4	15.4	57.3	15.4	1.7	0.0
Clay	31	*	*	*	*	*	*	*
Cleveland	762	0.3	11.7	17.1	58.9	11.9	0.1	0.0
Columbus	436	0.7	7.8	17.4	59.9	13.1	1.1	0.0
Craven	912	0.4	5.5	16.9	65.1	11.6	0.4	0.0
Cumberland	3,356	0.4	5.8	12.5	67.9	12.7	0.7	0.0
Currituck	84	*	*	*	*	*	*	*
Dare	180	0.6	3.9	7.2	66.7	20.0	1.7	0.0
Davidson	1,037	0.6	8.6	17.1	59.1	13.7	1.0	0.0
Davie	209	0.5	8.1	13.9	59.3	17.2	1.0	0.0
Duplin	594	0.2	6.9	14.0	59.6	18.4	1.0	0.0
Durham	1,884	0.4	6.8	11.5	58.6	21.7	1.1	0.0
Edgecombe	620	0.8	8.2	16.9	62.4	11.1	0.5	0.0
Forsyth	2,431	0.9	7.6	12.3	60.8	17.6	0.9	0.0
Franklin	316	0.0	8.5	19.0	50.3	21.8	0.3	0.0
Gaston	1,520	0.6	9.7	16.4	58.4	14.5	0.5	0.0
Gates	30	*	*	*	*	*	*	*
Graham	75	*	*	*	*	*	*	*

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

Table 4B
2007 Pregnancy Nutrition Surveillance ⁽¹⁾
North Carolina
Comparison of Age Distribution by County

Comparison of Age Distribution ⁽²⁾								
County	Number Included in Analysis	< 15 Years %	15 - 17 Years %	18 - 19 Years %	20 - 29 Years %	30 - 39 Years %	>= 40 Years %	Unknown %
Granville	325	1.2	6.2	10.8	63.1	17.8	0.9	0.0
Greene	221	0.5	10.4	17.6	58.8	11.8	0.9	0.0
Guilford	3,004	0.5	6.8	12.6	60.2	18.7	1.2	0.0
Halifax	551	0.5	8.2	16.9	58.1	15.1	1.3	0.0
Harnett	813	0.6	8.5	16.7	56.2	17.5	0.5	0.0
Haywood	318	0.9	8.5	16.7	58.2	15.1	0.6	0.0
Henderson	484	0.4	7.9	12.4	60.5	18.4	0.4	0.0
Hertford	204	1.5	8.8	16.7	63.7	8.3	1.0	0.0
Hoke	415	1.2	8.0	14.0	60.7	15.2	1.0	0.0
Hyde	30	*	*	*	*	*	*	*
Iredell	983	0.6	8.9	15.5	59.1	15.2	0.8	0.0
Jackson	255	0.4	7.8	13.7	59.6	18.4	0.0	0.0
Johnston	1,011	0.3	7.1	13.2	59.8	18.2	1.4	0.0
Jones	76	*	*	*	*	*	*	*
Lee	514	0.2	9.5	13.6	58.0	16.5	2.1	0.0
Lenoir	518	0.4	12.2	16.8	54.4	15.6	0.6	0.0
Lincoln	417	0.5	8.9	16.3	59.7	14.4	0.2	0.0
Macon	330	0.6	7.9	16.7	63.6	10.3	0.9	0.0
Madison	244	0.4	9.4	9.8	61.1	18.4	0.8	0.0

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

Table 4B
2007 Pregnancy Nutrition Surveillance ⁽¹⁾
North Carolina
Comparison of Age Distribution by County

Comparison of Age Distribution ⁽²⁾								
County	Number Included in Analysis	< 15 Years %	15 - 17 Years %	18 - 19 Years %	20 - 29 Years %	30 - 39 Years %	>= 40 Years %	Unknown %
Martin	117	0.0	9.4	13.7	65.0	12.0	0.0	0.0
Mcdowell	195	0.5	8.7	22.1	53.8	14.9	0.0	0.0
Mecklenbur	4,647	0.3	7.6	11.9	58.0	20.9	1.3	0.0
Mitchell	88	*	*	*	*	*	*	*
Montgomery	239	1.3	7.5	15.1	59.8	15.1	1.3	0.0
Moore	292	0.3	5.1	13.0	63.0	16.1	2.4	0.0
Nash	736	0.8	7.5	16.8	59.6	14.3	1.0	0.0
New Hanove	1,166	0.3	5.2	13.6	63.6	16.7	0.7	0.0
Northhampt	163	0.0	12.3	19.0	56.4	10.4	1.8	0.0
Onslow	2,134	0.1	3.4	14.2	70.9	10.8	0.4	0.0
Orange	490	0.4	5.7	8.4	64.1	20.0	1.4	0.0
Pamlico	73	*	*	*	*	*	*	*
Pasquotank	280	0.4	9.6	14.6	62.5	12.1	0.7	0.0
Pender	351	0.6	6.8	13.4	59.8	17.4	2.0	0.0
Perquimans	68	*	*	*	*	*	*	*
Person	272	0.4	10.7	15.4	59.9	13.2	0.4	0.0
Pitt	1,226	0.2	6.9	12.0	63.5	16.7	0.7	0.0
Polk	54	*	*	*	*	*	*	*
Randolph	998	0.6	9.3	14.1	58.2	17.0	0.7	0.0

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

Table 4B
2007 Pregnancy Nutrition Surveillance ⁽¹⁾
North Carolina
Comparison of Age Distribution by County

Comparison of Age Distribution ⁽²⁾								
County	Number Included in Analysis	< 15 Years %	15 - 17 Years %	18 - 19 Years %	20 - 29 Years %	30 - 39 Years %	>= 40 Years %	Unknown %
Richmond	528	1.3	11.7	14.6	57.0	14.4	0.9	0.0
Robeson	1,498	0.7	8.3	15.4	60.3	14.9	0.4	0.0
Rockingham	673	1.0	8.0	16.6	57.7	15.9	0.7	0.0
Rowan	977	0.5	9.6	14.7	60.5	13.9	0.7	0.0
Rutherford	395	0.3	6.8	15.7	62.3	13.9	1.0	0.0
Sampson	595	0.5	7.4	10.8	64.0	16.5	0.8	0.0
Scotland	443	0.7	12.6	15.1	58.5	11.7	1.4	0.0
Stanly	412	0.5	11.7	16.0	58.0	12.6	1.2	0.0
Stokes	252	0.0	7.5	11.1	63.9	17.1	0.4	0.0
Surry	509	0.2	8.3	17.5	54.4	17.9	1.8	0.0
Swain	155	0.0	9.0	12.3	65.2	12.3	1.3	0.0
Transylvan	169	0.0	8.3	21.3	56.2	13.0	1.2	0.0
Tyrrell	46	*	*	*	*	*	*	*
Union	1,014	0.8	8.0	13.1	57.3	19.3	1.5	0.0
Vance	496	0.6	10.5	16.7	56.3	15.5	0.4	0.0
Wake	4,418	0.3	6.3	11.0	59.1	22.1	1.1	0.0
Warren	167	0.6	15.6	11.4	57.5	13.8	1.2	0.0
Washington	147	0.0	11.6	15.0	57.8	14.3	1.4	0.0
Watauga	167	1.2	7.8	10.2	62.3	18.0	0.6	0.0

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

Table 4B
2007 Pregnancy Nutrition Surveillance ⁽¹⁾
North Carolina
Comparison of Age Distribution by County

Comparison of Age Distribution ⁽²⁾								
County	Number Included in Analysis	< 15 Years %	15 - 17 Years %	18 - 19 Years %	20 - 29 Years %	30 - 39 Years %	>= 40 Years %	Unknown %
Wayne Coun	1,088	0.4	7.7	13.4	62.6	15.0	0.9	0.0
Wilkes	505	0.2	9.5	14.5	60.4	14.9	0.6	0.0
Wilson Cou	737	1.2	9.9	13.8	58.1	16.0	0.9	0.0
Yadkin	264	0.4	12.1	10.6	57.6	18.6	0.8	0.0
Yancey	104	0.0	2.9	14.4	62.5	20.2	0.0	0.0
Unknown Counties	259	1.9	8.1	18.5	59.1	12.4	0.0	0.0
North Carolina	64,288	0.5	7.8	14.0	60.4	16.4	0.9	0.0
Nation (Prior Year)	1,153,475	0.4	6.5	12.6	60.0	19.0	1.4	0.1

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

Table 5B
2007 Pregnancy Nutrition Surveillance ⁽¹⁾
North Carolina
Comparison of Maternal Health Indicators by County

Comparison of Maternal Health Indicators ^(2, 3)																		
County	Prepregnancy BMI ⁽⁴⁾			Weight Gain ⁽⁵⁾			Anemia (Low Hb/Hct) ⁽⁶⁾				Gestational Diabetes During Pregnancy		Hypertension During Pregnancy					
	Under-weight	Over-weight		< Ideal	> Ideal		3rd Trimester		Postpartum									
	Number	% (Rank)	%	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)				
Alamance	676	10.5 (40)	45.0 (26)	340	24.4 (21)	38.5 (30)	127	33.9 (10)	626	35.6 (38)	0	*	*	0	*	*		
Alexander	202	12.9 (64)	44.6 (24)	157	30.6 (53)	35.0 (18)	44	*	*	87	*	*	0	*	*	0	*	*
Alleghany	45	*	*	35	*	*	7	*	*	20	*	*	0	*	*	0	*	*
Anson	192	13.0 (67)	50.0 (57)	165	27.3 (41)	41.8 (49)	24	*	*	42	*	*	0	*	*	0	*	*
Ashe	144	15.3 (80)	44.4 (20)	125	27.2 (40)	37.6 (26)	23	*	*	34	*	*	0	*	*	0	*	*
Avery	102	17.6 (86)	40.2 (3)	11	*	*	12	*	*	20	*	*	0	*	*	0	*	*
Beaufort	301	9.0 (23)	57.8 (82)	140	25.0 (25)	32.9 (12)	34	*	*	249	38.2 (42)	0	*	*	0	*	*	
Bertie	148	6.1 (4)	67.6 (86)	82	*	*	26	*	*	81	*	*	0	*	*	0	*	*
Bladen	283	8.1 (14)	50.5 (61)	137	32.8 (62)	35.8 (21)	18	*	*	64	*	*	0	*	*	0	*	*
Brunswick	160	16.3 (84)	43.8 (16)	71	*	*	5	*	*	201	13.9 (2)	0	*	*	0	*	*	
Buncombe	1,209	13.9 (75)	42.1 (9)	766	19.1 (5)	43.5 (55)	194	25.8 (7)	480	34.2 (37)	0	*	*	0	*	*		
Burke	499	11.2 (47)	44.7 (25)	396	27.5 (42)	31.8 (6)	116	34.5 (11)	254	31.5 (32)	0	*	*	0	*	*		
Cabarrus	1,007	10.3 (36)	42.8 (10)	853	29.1 (45)	38.8 (32)	107	34.6 (12)	239	22.6 (16)	0	*	*	0	*	*		
Caldwell	443	13.8 (74)	46.5 (31)	367	24.0 (19)	46.3 (65)	50	*	*	194	16.0 (3)	0	*	*	0	*	*	
Camden	23	*	*	13	*	*	2	*	*	18	*	*	0	*	*	0	*	*
Carteret	261	11.1 (45)	41.4 (5)	228	14.0 (1)	46.1 (64)	79	*	*	137	20.4 (11)	0	*	*	0	*	*	
Caswell	94	*	*	75	*	*	11	*	*	25	*	*	0	*	*	0	*	*

- (1) Reporting period is January 1 through December 31.
- (2) Excludes records with unknown data or errors.
- (3) Rank compares one county's rate to other counties. Rank 1 = best rate.
- (4) Based on 1990 IOM report, "Nutrition During Pregnancy": underweight (BMI <19.8), overweight (BMI >26.0; includes overweight and obese women).
- (5) Based on 1990 IOM report, "Nutrition During Pregnancy": underweight prepregnancy (ideal weight gain = 28 to 40 lbs), normal weight prepregnancy (ideal weight gain = 25 to 35 lbs), overweight prepregnancy (ideal weight gain = 15 to 25 lbs), obese prepregnancy (ideal weight gain = 15 to 25 lbs).
- (6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.
- * Percentages are not calculated if <100 records are available for analysis after exclusions.

Table 5B
2007 Pregnancy Nutrition Surveillance ⁽¹⁾
North Carolina
Comparison of Maternal Health Indicators by County

Comparison of Maternal Health Indicators ^(2, 3)														
County	Prepregnancy BMI (4)			Weight Gain (5)			Anemia (Low Hb/Hct) (6)				Gestational Diabetes During Pregnancy		Hypertension During Pregnancy	
	Under- weight		Over- weight	< Ideal		> Ideal	3rd Trimester		Postpartum					
	Number	% (Rank)	%	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Catawba	954	9.9 (30)	46.9 (33)	738	38.5 (70)	30.6 (4)	197	25.4 (5)	622	25.2(22)	0	* *	0	* *
Chatham	257	8.2 (15)	50.2 (59)	206	26.7 (35)	32.0 (8)	19	* *	230	23.0(18)	0	* *	0	* *
Cherokee	117	11.1 (45)	48.7 (48)	91	* *	* *	30	* *	34	* *	0	* *	0	* *
Chowan	104	8.7 (20)	60.6 (84)	76	* *	* *	15	* *	93	* *	0	* *	0	* *
Clay	29	* *	* *	21	* *	* *	5	* *	12	* *	0	* *	0	* *
Cleveland	581	12.7 (60)	45.8 (29)	22	* *	* *	68	* *	544	28.5(25)	0	* *	0	* *
Columbus	356	10.4 (39)	50.3 (60)	147	25.9 (27)	41.5 (47)	66	* *	51	* *	0	* *	0	* *
Craven	590	11.5 (51)	44.2 (19)	406	23.2 (17)	36.9 (25)	29	* *	584	23.1(19)	0	* *	0	* *
Cumberland	2,532	12.3 (58)	43.3 (14)	1,591	22.9 (16)	47.3 (68)	572	50.9 (21)	806	41.4(46)	0	* *	0	* *
Currituck	66	* *	* *	49	* *	* *	15	* *	66	* *	0	* *	0	* *
Dare	149	13.4 (70)	43.0 (11)	116	27.6 (43)	34.5 (16)	12	* *	67	* *	0	* *	0	* *
Davidson	760	13.0 (67)	41.6 (7)	601	27.1 (39)	36.3 (24)	124	19.4 (3)	451	17.1 (6)	0	* *	0	* *
Davie	173	14.5 (78)	44.5 (22)	141	22.7 (15)	44.7 (61)	30	* *	112	16.1 (5)	0	* *	0	* *
Duplin	288	7.6 (11)	57.3 (81)	221	31.7 (59)	34.4 (15)	75	* *	176	43.8(48)	0	* *	0	* *
Durham	1,242	9.9 (30)	46.0 (30)	1,032	31.4 (58)	35.4 (20)	41	* *	119	28.6(27)	0	* *	0	* *
Edgecombe	247	7.3 (8)	66.0 (85)	86	* *	* *	56	* *	271	38.7(43)	0	* *	0	* *
Forsyth	1,710	11.3 (49)	44.0 (18)	1,158	26.2 (32)	40.7 (42)	237	38.4 (16)	829	35.8(39)	0	* *	0	* *

- (1) Reporting period is January 1 through December 31.
- (2) Excludes records with unknown data or errors.
- (3) Rank compares one county's rate to other counties. Rank 1 = best rate.
- (4) Based on 1990 IOM report, "Nutrition During Pregnancy": underweight (BMI <19.8), overweight (BMI >26.0; includes overweight and obese women).
- (5) Based on 1990 IOM report, "Nutrition During Pregnancy": underweight prepregnancy (ideal weight gain = 28 to 40 lbs), normal weight prepregnancy (ideal weight gain = 25 to 35 lbs), overweight prepregnancy (ideal weight gain = 15 to 25 lbs), obese prepregnancy (ideal weight gain = 15 to 25 lbs).
- (6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.
- * Percentages are not calculated if <100 records are available for analysis after exclusions.

Table 5B
2007 Pregnancy Nutrition Surveillance ⁽¹⁾
North Carolina
Comparison of Maternal Health Indicators by County

Comparison of Maternal Health Indicators ^(2, 3)														
County	Prepregnancy BMI ⁽⁴⁾			Weight Gain ⁽⁵⁾			Anemia (Low Hb/Hct) ⁽⁶⁾				Gestational Diabetes During Pregnancy	Hypertension During Pregnancy		
	Under-weight		Over-weight	< Ideal		> Ideal	3rd Trimester		Postpartum					
	Number	% (Rank)	%	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)			Number	% (Rank)
Franklin	262	6.9 (6)	51.5 (68)	194	32.5 (60)	32.0 (8)	44	* *	47	* *	0	* *	0	* *
Gaston	1,237	15.6 (82)	44.5 (22)	978	31.3 (57)	39.5 (39)	268	23.1 (4)	699	50.5 (51)	0	* *	0	* *
Gates	24	* *	* *	19	* *	* *	2	* *	18	* *	0	* *	0	* *
Graham	64	* *	* *	52	* *	* *	2	* *	21	* *	0	* *	0	* *
Granville	259	6.9 (6)	53.3 (71)	211	29.4 (47)	38.9 (34)	30	* *	78	* *	0	* *	0	* *
Greene	176	9.1 (25)	50.6 (63)	153	34.6 (66)	34.0 (13)	22	* *	68	* *	0	* *	0	* *
Guilford	2,238	10.3 (36)	46.9 (33)	1,049	26.6 (34)	40.7 (42)	169	32.0 (9)	639	22.4 (15)	0	* *	0	* *
Halifax	436	7.3 (8)	55.3 (74)	391	25.6 (26)	42.7 (53)	41	* *	204	52.9 (52)	0	* *	0	* *
Harnett	657	10.0 (33)	48.7 (48)	464	24.1 (20)	43.8 (57)	143	39.2 (18)	277	32.5 (33)	0	* *	0	* *
Haywood	254	11.8 (52)	40.6 (4)	161	17.4 (3)	42.2 (50)	25	* *	131	16.0 (3)	0	* *	0	* *
Henderson	352	10.2 (35)	47.4 (38)	177	22.0 (11)	45.2 (63)	97	* *	214	33.6 (36)	0	* *	0	* *
Hertford	176	8.0 (13)	60.2 (83)	117	26.5 (33)	38.5 (30)	13	* *	51	* *	0	* *	0	* *
Hoke	135	7.4 (10)	46.7 (32)	115	23.5 (18)	40.0 (41)	11	* *	76	* *	0	* *	0	* *
Hyde	25	* *	* *	20	* *	* *	7	* *	8	* *	0	* *	0	* *
Iredell	803	12.8 (62)	46.9 (33)	721	26.9 (36)	42.3 (52)	175	26.9 (8)	804	13.7 (1)	0	* *	0	* *
Jackson	190	8.9 (22)	47.4 (38)	148	20.3 (7)	41.2 (45)	24	* *	121	22.3 (14)	0	* *	0	* *
Johnston	567	10.1 (34)	48.5 (46)	301	30.9 (54)	34.6 (17)	150	41.3 (19)	544	31.4 (31)	0	* *	0	* *

- (1) Reporting period is January 1 through December 31.
- (2) Excludes records with unknown data or errors.
- (3) Rank compares one county's rate to other counties. Rank 1 = best rate.
- (4) Based on 1990 IOM report, "Nutrition During Pregnancy": underweight (BMI <19.8), overweight (BMI >26.0; includes overweight and obese women).
- (5) Based on 1990 IOM report, "Nutrition During Pregnancy": underweight prepregnancy (ideal weight gain = 28 to 40 lbs), normal weight prepregnancy (ideal weight gain = 25 to 35 lbs), overweight prepregnancy (ideal weight gain = 15 to 25 lbs), obese prepregnancy (ideal weight gain = 15 to 25 lbs).
- (6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.
- * Percentages are not calculated if <100 records are available for analysis after exclusions.

Table 5B
2007 Pregnancy Nutrition Surveillance ⁽¹⁾
North Carolina
Comparison of Maternal Health Indicators by County

Comparison of Maternal Health Indicators ^(2, 3)																		
County	Prepregnancy BMI ⁽⁴⁾				Weight Gain ⁽⁵⁾			Anemia (Low Hb/Hct) ⁽⁶⁾				Gestational Diabetes During Pregnancy	Hypertension During Pregnancy					
	Under-weight		Over-weight		< Ideal		> Ideal	3rd Trimester		Postpartum								
	Number	% (Rank)	%	%	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)			Number	% (Rank)			
Jones	68	* * * *			5	* * * *			11	* * *			6	* * *	0	* * *	0	* * *
Lee	401	6.0 (3)	47.9 (41)		315	24.4 (21)	42.2 (50)		54	* * *			192	32.8 (34)	0	* * *	0	* * *
Lenoir	223	12.1 (56)	43.0 (11)		146	26.0 (29)	44.5 (59)		85	* * *			298	40.3 (44)	0	* * *	0	* * *
Lincoln	331	13.6 (72)	49.5 (55)		227	22.5 (14)	44.1 (58)		36	* * *			164	17.1 (6)	0	* * *	0	* * *
Macon	252	11.9 (53)	49.2 (54)		195	34.4 (65)	30.3 (3)		35	* * *			116	25.0 (21)	0	* * *	0	* * *
Madison	214	12.6 (59)	43.5 (15)		109	20.2 (6)	39.4 (37)		39	* * *			80	* * *	0	* * *	0	* * *
Martin	101	12.9 (64)	52.5 (69)		21	* * * *			12	* * *			19	* * *	0	* * *	0	* * *
Mcdowell	165	12.7 (60)	55.8 (77)		132	29.5 (48)	39.4 (37)		35	* * *			40	* * *	0	* * *	0	* * *
Mecklenbur	2,219	10.6 (41)	48.2 (42)		1,467	24.8 (24)	41.4 (46)		485	36.3 (15)			1,786	35.8 (39)	0	* * *	0	* * *
Mitchell	74	* * * *			59	* * * *			7	* * *			33	* * *	0	* * *	0	* * *
Montgomery	171	8.2 (15)	53.2 (70)		53	* * * *			17	* * *			32	* * *	0	* * *	0	* * *
Moore	173	5.2 (1)	56.1 (78)		51	* * * *			29	* * *			61	* * *	0	* * *	0	* * *
Nash	594	10.3 (36)	50.0 (57)		453	32.9 (64)	34.2 (14)		55	* * *			142	42.3 (47)	0	* * *	0	* * *
New Hanove	672	11.2 (47)	48.7 (48)		387	22.2 (13)	41.6 (48)		81	* * *			398	33.4 (35)	0	* * *	0	* * *
Northhampt	135	11.9 (53)	55.6 (75)		83	* * * *			22	* * *			42	* * *	0	* * *	0	* * *
Onslow	1,545	13.7 (73)	41.8 (8)		1,262	18.5 (4)	46.7 (66)		327	25.7 (6)			1,348	22.2 (13)	0	* * *	0	* * *
Orange	308	7.8 (12)	48.4 (45)		225	24.4 (21)	43.6 (56)		11	* * *			198	28.8 (28)	0	* * *	0	* * *

- (1) Reporting period is January 1 through December 31.
- (2) Excludes records with unknown data or errors.
- (3) Rank compares one county's rate to other counties. Rank 1 = best rate.
- (4) Based on 1990 IOM report, "Nutrition During Pregnancy": underweight (BMI <19.8), overweight (BMI >26.0; includes overweight and obese women).
- (5) Based on 1990 IOM report, "Nutrition During Pregnancy": underweight prepregnancy (ideal weight gain = 28 to 40 lbs), normal weight prepregnancy (ideal weight gain = 25 to 35 lbs), overweight prepregnancy (ideal weight gain = 15 to 25 lbs), obese prepregnancy (ideal weight gain = 15 to 25 lbs).
- (6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.
- * Percentages are not calculated if <100 records are available for analysis after exclusions.

Table 5B
2007 Pregnancy Nutrition Surveillance ⁽¹⁾
North Carolina
Comparison of Maternal Health Indicators by County

Comparison of Maternal Health Indicators ^(2, 3)														
County	Prepregnancy BMI ⁽⁴⁾				Weight Gain ⁽⁵⁾				Anemia (Low Hb/Hct) ⁽⁶⁾				Gestational Diabetes During Pregnancy	Hypertension During Pregnancy
	Under-weight		Over-weight		< Ideal		> Ideal		3rd Trimester		Postpartum			
	Number	% (Rank)	%	%	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number		
Pamlico	61	* * * *			7	* * * *			2	* * *	9	* * *	0	* * *
Pasquotank	195	9.2 (26)	56.9 (80)		150	36.0 (69)	31.3 (5)		37	* * *	237	75.5 (54)	0	* * *
Pender	263	9.9 (30)	48.3 (44)		230	29.1 (45)	36.1 (23)		53	* * *	67	* * *	0	* * *
Perquimans	41	* * * *			4	* * * *			9	* * *	45	* * *	0	* * *
Person	228	11.0 (43)	48.7 (48)		120	14.2 (2)	49.2 (70)		33	* * *	93	* * *	0	* * *
Pitt	994	8.5 (19)	54.1 (72)		766	27.0 (37)	38.9 (34)		178	57.9 (22)	215	48.8 (50)	0	* * *
Polk	41	* * * *			18	* * * *			7	* * *	14	* * *	0	* * *
Randolph	768	13.2 (69)	49.1 (52)		601	21.5 (9)	44.6 (60)		171	19.3 (2)	670	20.3 (10)	0	* * *
Richmond	395	17.2 (85)	41.5 (6)		287	26.1 (30)	43.2 (54)		81	* * *	181	22.7 (17)	0	* * *
Robeson	1,204	13.9 (75)	47.6 (40)		793	30.4 (51)	35.3 (19)		270	16.3 (1)	407	29.7 (30)	0	* * *
Rockingham	281	10.7 (42)	43.8 (16)		75	* * * *			75	* * *	170	17.1 (6)	0	* * *
Rowan	785	13.9 (75)	45.5 (28)		622	27.0 (37)	39.7 (40)		194	38.7 (17)	358	36.9 (41)	0	* * *
Rutherford	266	15.0 (79)	45.1 (27)		112	41.1 (71)	29.5 (2)		28	* * *	193	29.5 (29)	0	* * *
Sampson	407	5.7 (2)	56.3 (79)		310	20.6 (8)	47.4 (69)		80	* * *	230	27.4 (23)	0	* * *
Scotland	264	6.4 (5)	54.9 (73)		180	32.8 (62)	41.1 (44)		22	* * *	58	* * *	0	* * *
Stanly	345	15.7 (83)	39.1 (2)		234	34.6 (66)	38.9 (34)		49	* * *	231	48.1 (49)	0	* * *
Stokes	163	12.9 (64)	48.5 (46)		119	26.1 (30)	51.3 (71)		33	* * *	51	* * *	0	* * *

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one county's rate to other counties. Rank 1 = best rate.
(4) Based on 1990 IOM report, "Nutrition During Pregnancy": underweight (BMI <19.8), overweight (BMI >26.0; includes overweight and obese women).
(5) Based on 1990 IOM report, "Nutrition During Pregnancy": underweight prepregnancy (ideal weight gain = 28 to 40 lbs), normal weight prepregnancy (ideal weight gain = 25 to 35 lbs), overweight prepregnancy (ideal weight gain = 15 to 25 lbs), obese prepregnancy (ideal weight gain = 15 to 25 lbs).
(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

Table 5B
2007 Pregnancy Nutrition Surveillance ⁽¹⁾
North Carolina
Comparison of Maternal Health Indicators by County

Comparison of Maternal Health Indicators ^(2, 3)														
County	Prepregnancy BMI (4)			Weight Gain (5)			Anemia (Low Hb/Hct) (6)				Gestational Diabetes During Pregnancy		Hypertension During Pregnancy	
	Under-weight		Over-weight	< Ideal		> Ideal	3rd Trimester		Postpartum					
	Number	% (Rank)	%	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Surry	335	13.4 (70)	43.0 (11)	235	22.1 (12)	47.2 (67)	64	* *	155	41.3 (45)	0	* *	0	* *
Swain	112	9.8 (29)	48.2 (42)	31	* *	* *	13	* *	44	* *	0	* *	0	* *
Transylvan	136	15.4 (81)	35.3 (1)	59	* *	* *	17	* *	30	* *	0	* *	0	* *
Tyrrell	39	* *	* *	26	* *	* *	3	* *	9	* *	0	* *	0	* *
Union	708	11.3 (49)	47.0 (36)	405	30.4 (51)	32.3 (10)	153	35.9 (14)	426	23.5 (20)	0	* *	0	* *
Vance	335	11.0 (43)	51.0 (66)	184	31.0 (56)	35.9 (22)	27	* *	130	57.7 (53)	0	* *	0	* *
Wake	2,564	9.0 (23)	47.2 (37)	1,533	32.6 (61)	32.4 (11)	282	34.8 (13)	1,375	21.5 (12)	0	* *	0	* *
Warren	144	9.7 (27)	51.4 (67)	104	28.8 (44)	28.8 (1)	15	* *	46	* *	0	* *	0	* *
Washington	133	8.3 (17)	55.6 (75)	103	30.1 (50)	37.9 (28)	16	* *	16	* *	0	* *	0	* *
Watauga	126	11.9 (53)	44.4 (20)	50	* *	* *	10	* *	63	* *	0	* *	0	* *
Wayne Coun	770	8.8 (21)	50.5 (61)	398	30.9 (54)	37.9 (28)	67	* *	400	28.5 (25)	0	* *	0	* *
Wilkes	456	12.1 (56)	49.1 (52)	337	35.3 (68)	31.8 (6)	88	* *	148	18.2 (9)	0	* *	0	* *
Wilson Cou	649	9.7 (27)	50.8 (65)	449	29.8 (49)	37.6 (26)	132	44.7 (20)	320	27.8 (24)	0	* *	0	* *
Yadkin	225	8.4 (18)	50.7 (64)	190	21.6 (10)	44.7 (61)	34	* *	89	* *	0	* *	0	* *
Yancey	76	* *	* *	29	* *	* *	4	* *	32	* *	0	* *	0	* *
Unknown Counties	203	12.8 (62)	49.8 (56)	139	25.9 (27)	38.8 (32)	20	* *	95	* *	0	* *	0	* *
North Carolina	45,157	11.2	47.2	30,024	27.0	39.5	7,301	34.7	23,400	30.8	0	*	0	*

- (1) Reporting period is January 1 through December 31.
 - (2) Excludes records with unknown data or errors.
 - (3) Rank compares one county's rate to other counties. Rank 1 = best rate.
 - (4) Based on 1990 IOM report, "Nutrition During Pregnancy": underweight (BMI <19.8), overweight (BMI >26.0; includes overweight and obese women).
 - (5) Based on 1990 IOM report, "Nutrition During Pregnancy": underweight prepregnancy (ideal weight gain = 28 to 40 lbs), normal weight prepregnancy (ideal weight gain = 25 to 35 lbs), overweight prepregnancy (ideal weight gain = 15 to 25 lbs), obese prepregnancy (ideal weight gain = 15 to 25 lbs).
 - (6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.
- * Percentages are not calculated if <100 records are available for analysis after exclusions.

Table 5B
2007 Pregnancy Nutrition Surveillance ⁽¹⁾
North Carolina
Comparison of Maternal Health Indicators by County

Comparison of Maternal Health Indicators ^(2, 3)														
County	Prepregnancy BMI (4)			Weight Gain (5)			Anemia (Low Hb/Hct) (6)				Gestational Diabetes During Pregnancy		Hypertension During Pregnancy	
	Under- weight		Over- weight	< Ideal		> Ideal	3rd Trimester		Postpartum					
	Number	% (Rank)	%	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Nation (Prior Year)	1,081,550	11.2	43.7	872,976	24.8	43.1	161,665	32.2	359,221	29.2	150,176	3.6	164,043	3.0

- (1) Reporting period is January 1 through December 31.
- (2) Excludes records with unknown data or errors.
- (3) Rank compares one county's rate to other counties. Rank 1 = best rate.
- (4) Based on 1990 IOM report, "Nutrition During Pregnancy": underweight (BMI <19.8), overweight (BMI >26.0; includes overweight and obese women).
- (5) Based on 1990 IOM report, "Nutrition During Pregnancy": underweight prepregnancy (ideal weight gain = 28 to 40 lbs), normal weight prepregnancy (ideal weight gain = 25 to 35 lbs), overweight prepregnancy (ideal weight gain = 15 to 25 lbs), obese prepregnancy (ideal weight gain = 15 to 25 lbs).
- (6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.
- * Percentages are not calculated if <100 records are available for analysis after exclusions.

Table 6B
2007 Pregnancy Nutrition Surveillance ⁽¹⁾
North Carolina
Comparison of Maternal Behavioral Indicators by County

Comparison of Maternal Behavioral Indicators ^(2, 3)										
County	Medical Care			WIC Enrollment		Multivitamin Consumption				
	1st Trimester		No Care	1st Trimester		Prepregnancy (4)		During Pregnancy (5)		
	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	
Alamance	1,017	71.2 (64)	0.5 (21)	1,142	39.1 (26)	0	* *	0	* *	
Alexander	203	71.4 (62)	2.5 (82)	234	28.6 (53)	0	* *	0	* *	
Alleghany	46	* *	* *	54	* *	0	* *	0	* *	
Anson	210	74.8 (47)	0.0 (1)	227	41.0 (19)	0	* *	0	* *	
Ashe	154	85.7 (13)	1.3 (69)	160	47.5 (4)	0	* *	0	* *	
Avery	29	* *	* *	119	41.2 (17)	0	* *	0	* *	
Beaufort	288	73.3 (56)	1.4 (70)	370	45.4 (6)	0	* *	0	* *	
Bertie	116	84.5 (16)	0.0 (1)	171	39.2 (25)	0	* *	0	* *	
Bladen	171	70.2 (65)	1.2 (64)	303	43.9 (13)	0	* *	0	* *	
Brunswick	439	79.7 (34)	0.5 (21)	611	33.2 (37)	0	* *	0	* *	
Buncombe	1,135	92.3 (1)	0.4 (17)	1,410	28.3 (54)	0	* *	0	* *	
Burke	586	57.8 (82)	0.5 (21)	583	17.7 (82)	0	* *	0	* *	
Cabarrus	1,189	63.7 (75)	0.8 (40)	1,181	24.6 (66)	0	* *	0	* *	
Caldwell	498	91.6 (2)	0.2 (11)	522	37.4 (28)	0	* *	0	* *	
Camden	26	* *	* *	29	* *	0	* *	0	* *	
Carteret	318	83.6 (19)	0.6 (30)	336	21.1 (76)	0	* *	0	* *	
Caswell	104	78.8 (36)	0.0 (1)	116	44.8 (9)	0	* *	0	* *	
Catawba	871	75.9 (42)	0.7 (35)	1,000	22.5 (70)	0	* *	0	* *	
Chatham	303	73.6 (55)	0.7 (35)	333	33.6 (36)	0	* *	0	* *	
Cherokee	120	71.7 (60)	0.8 (40)	137	29.2 (52)	0	* *	0	* *	

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one county's rate to other counties. Rank 1 = best rate.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

Table 6B
2007 Pregnancy Nutrition Surveillance ⁽¹⁾
North Carolina
Comparison of Maternal Behavioral Indicators by County

Comparison of Maternal Behavioral Indicators ^(2, 3)										
County	Medical Care				WIC Enrollment		Multivitamin Consumption			
	1st Trimester		No Care		1st Trimester		Prepregnancy (4)		During Pregnancy (5)	
	Number	% (Rank)	% (Rank)		Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Chowan	105	88.6 (8)	0.0 (1)		112	41.1 (18)	0	* *	0	* *
Clay	26	* *	* *		30	* *	0	* *	0	* *
Cleveland	585	68.5 (67)	0.7 (35)		733	40.5 (21)	0	* *	0	* *
Columbus	220	59.5 (80)	0.5 (21)		415	32.8 (40)	0	* *	0	* *
Craven	748	79.9 (33)	0.1 (10)		887	29.8 (50)	0	* *	0	* *
Cumberland	2,984	83.4 (21)	1.0 (54)		3,252	22.4 (71)	0	* *	0	* *
Currituck	75	* *	* *		79	* *	0	* *	0	* *
Dare	162	81.5 (28)	0.0 (1)		176	36.9 (30)	0	* *	0	* *
Davidson	934	74.4 (48)	0.4 (17)		992	27.1 (59)	0	* *	0	* *
Davie	192	88.0 (10)	1.0 (54)		202	31.7 (42)	0	* *	0	* *
Duplin	571	75.1 (46)	0.9 (50)		584	26.7 (60)	0	* *	0	* *
Durham	1,726	81.6 (27)	0.8 (40)		1,801	10.1 (88)	0	* *	0	* *
Edgecombe	565	62.1 (79)	2.3 (80)		579	27.8 (56)	0	* *	0	* *
Forsyth	2,116	81.0 (31)	0.5 (21)		2,347	17.7 (82)	0	* *	0	* *
Franklin	274	81.8 (26)	0.4 (17)		299	26.1 (61)	0	* *	0	* *
Gaston	1,346	51.9 (83)	0.9 (50)		1,429	20.7 (77)	0	* *	0	* *
Gates	30	* *	* *		28	* *	0	* *	0	* *
Graham	64	* *	* *		72	* *	0	* *	0	* *
Granville	295	75.6 (43)	1.0 (54)		307	27.7 (57)	0	* *	0	* *
Greene	209	84.2 (17)	1.0 (54)		206	33.0 (38)	0	* *	0	* *

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one county's rate to other counties. Rank 1 = best rate.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

Table 6B
2007 Pregnancy Nutrition Surveillance ⁽¹⁾
North Carolina
Comparison of Maternal Behavioral Indicators by County

Comparison of Maternal Behavioral Indicators ^(2, 3)										
County	Medical Care			WIC Enrollment		Multivitamin Consumption				
	1st Trimester		No Care	1st Trimester		Prepregnancy (4)		During Pregnancy (5)		
	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	
Guilford	2,486	72.9 (58)	1.2 (64)	2,926	21.9 (73)	0	* *	0	* *	
Halifax	509	84.9 (14)	1.0 (54)	534	24.7 (65)	0	* *	0	* *	
Harnett	667	68.1 (69)	1.0 (54)	791	29.7 (51)	0	* *	0	* *	
Haywood	256	90.2 (4)	0.0 (1)	308	39.6 (23)	0	* *	0	* *	
Henderson	346	90.2 (4)	0.6 (30)	462	18.0 (81)	0	* *	0	* *	
Hertford	146	83.6 (19)	0.0 (1)	196	31.6 (43)	0	* *	0	* *	
Hoke	216	78.2 (40)	2.3 (80)	405	23.5 (68)	0	* *	0	* *	
Hyde	26	* *	* *	29	* *	0	* *	0	* *	
Iredell	912	58.9 (81)	2.0 (79)	956	23.4 (69)	0	* *	0	* *	
Jackson	225	90.7 (3)	0.0 (1)	249	45.0 (8)	0	* *	0	* *	
Johnston	887	78.8 (36)	0.6 (30)	991	16.3 (84)	0	* *	0	* *	
Jones	12	* *	* *	74	* *	0	* *	0	* *	
Lee	443	63.2 (77)	0.2 (11)	495	20.6 (78)	0	* *	0	* *	
Lenoir	388	62.6 (78)	0.5 (21)	486	27.6 (58)	0	* *	0	* *	
Lincoln	317	68.5 (67)	0.6 (30)	399	36.8 (31)	0	* *	0	* *	
Macon	291	83.2 (22)	0.0 (1)	311	42.4 (16)	0	* *	0	* *	
Madison	136	83.8 (18)	0.7 (35)	233	31.3 (45)	0	* *	0	* *	
Martin	37	* *	* *	115	45.2 (7)	0	* *	0	* *	
Mcdowell	170	78.8 (36)	1.2 (64)	185	35.7 (33)	0	* *	0	* *	
Mecklenbur	3,612	65.8 (72)	1.9 (77)	4,562	10.8 (87)	0	* *	0	* *	

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one county's rate to other counties. Rank 1 = best rate.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

Table 6B
2007 Pregnancy Nutrition Surveillance ⁽¹⁾
North Carolina
Comparison of Maternal Behavioral Indicators by County

Comparison of Maternal Behavioral Indicators ^(2, 3)										
County	Medical Care				WIC Enrollment		Multivitamin Consumption			
	1st Trimester		No Care		1st Trimester		Pregpregnancy (4)		During Pregpregnancy (5)	
	Number	% (Rank)	%	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Mitchell	76	* *	* *	* *	80	* *	0	* *	0	* *
Montgomery	110	75.5 (44)	3.6 (83)		229	43.7 (14)	0	* *	0	* *
Moore	193	81.9 (25)	1.0 (54)		285	21.8 (75)	0	* *	0	* *
Nash	649	68.1 (69)	1.4 (70)		693	30.9 (46)	0	* *	0	* *
New Hanove	954	74.1 (49)	1.9 (77)		1,148	31.6 (43)	0	* *	0	* *
Northhampt	120	82.5 (24)	0.8 (40)		155	21.9 (73)	0	* *	0	* *
Onslow	1,777	88.2 (9)	0.2 (11)		2,077	26.1 (61)	0	* *	0	* *
Orange	397	73.0 (57)	0.8 (40)		467	22.3 (72)	0	* *	0	* *
Pamlico	17	* *	* *	* *	71	* *	0	* *	0	* *
Pasquotank	263	74.1 (49)	0.8 (40)		271	25.8 (63)	0	* *	0	* *
Pender	331	71.9 (59)	1.2 (64)		343	28.3 (54)	0	* *	0	* *
Perquimans	65	* *	* *	* *	67	* *	0	* *	0	* *
Person	168	86.3 (12)	1.8 (76)		265	37.4 (28)	0	* *	0	* *
Pitt	1,047	84.7 (15)	0.5 (21)		1,179	32.3 (41)	0	* *	0	* *
Polk	28	* *	* *	* *	54	* *	0	* *	0	* *
Randolph	879	75.4 (45)	0.6 (30)		964	25.8 (63)	0	* *	0	* *
Richmond	449	71.5 (61)	1.6 (74)		511	30.9 (46)	0	* *	0	* *
Robeson	1,134	66.7 (71)	0.4 (17)		1,419	20.4 (79)	0	* *	0	* *
Rockingham	605	81.2 (29)	0.3 (14)		659	44.0 (11)	0	* *	0	* *
Rowan	883	63.6 (76)	1.1 (62)		940	15.7 (85)	0	* *	0	* *

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one county's rate to other counties. Rank 1 = best rate.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

Table 6B
2007 Pregnancy Nutrition Surveillance ⁽¹⁾
North Carolina
Comparison of Maternal Behavioral Indicators by County

Comparison of Maternal Behavioral Indicators ^(2, 3)										
County	Medical Care			WIC Enrollment		Multivitamin Consumption				
	1st Trimester		No Care	1st Trimester		Prepregnancy (4)		During Pregnancy (5)		
	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	
Rutherford	287	82.6 (23)	0.3 (14)	379	35.9 (32)	0	* *	0	* *	
Sampson	543	73.8 (53)	0.7 (35)	571	30.6 (48)	0	* *	0	* *	
Scotland	353	73.9 (52)	1.4 (70)	427	49.6 (3)	0	* *	0	* *	
Stanly	309	73.8 (53)	0.3 (14)	401	35.7 (33)	0	* *	0	* *	
Stokes	215	87.4 (11)	0.9 (50)	241	44.0 (11)	0	* *	0	* *	
Surry	411	80.3 (32)	0.5 (21)	495	38.6 (27)	0	* *	0	* *	
Swain	81	* *	* *	149	55.7 (1)	0	* *	0	* *	
Transylvan	91	* *	* *	167	55.1 (2)	0	* *	0	* *	
Tyrrell	36	* *	* *	43	* *	0	* *	0	* *	
Union	780	71.4 (62)	1.4 (70)	985	12.3 (86)	0	* *	0	* *	
Vance	399	68.9 (66)	0.8 (40)	470	23.6 (67)	0	* *	0	* *	
Wake	3,948	64.0 (74)	0.9 (50)	4,204	20.1 (80)	0	* *	0	* *	
Warren	127	74.0 (51)	0.8 (40)	160	42.5 (15)	0	* *	0	* *	
Washington	121	79.3 (35)	1.7 (75)	143	45.5 (5)	0	* *	0	* *	
Watauga	101	89.1 (6)	1.0 (54)	163	44.2 (10)	0	* *	0	* *	
Wayne Coun	682	65.7 (73)	1.2 (64)	1,060	34.6 (35)	0	* *	0	* *	
Wilkes	398	81.2 (29)	0.8 (40)	497	33.0 (38)	0	* *	0	* *	
Wilson Cou	622	78.3 (39)	1.1 (62)	705	30.4 (49)	0	* *	0	* *	
Yadkin	248	88.7 (7)	0.8 (40)	260	39.6 (23)	0	* *	0	* *	
Yancey	49	* *	* *	102	40.2 (22)	0	* *	0	* *	

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one county's rate to other counties. Rank 1 = best rate.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

Table 6B
2007 Pregnancy Nutrition Surveillance ⁽¹⁾
North Carolina
Comparison of Maternal Behavioral Indicators by County

Comparison of Maternal Behavioral Indicators ^(2, 3)									
County	Medical Care			WIC Enrollment		Multivitamin Consumption			
	1st Trimester		No Care	1st Trimester		Pregnancy (4)		During Pregnancy (5)	
	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Unknown Counties	212	76.9 (41)	0.5 (21)	249	41.0 (19)	0	* *	0	* *
North Carolina	53,290	74.3	0.9	62,023	25.9	0	*	0	*
Nation (Prior Year)	579,816	78.5	5.6	944,971	31.2	38,592	15.7	206,967	74.1

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one county's rate to other counties. Rank 1 = best rate.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

Table 7B
2007 Pregnancy Nutrition Surveillance (1)
North Carolina
Comparison of Smoking Indicators by County

Comparison of Smoking Indicators (2, 3)										
County	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)		Smoking in Household			
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy				Prenatal		Postpartum	
	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Alamance	986	33.2 (58)	1,006	21.7 (59)	271	31.4 (15)	0	* *	0	* *
Alexander	201	35.3 (68)	203	28.1 (78)	53	* *	0	* *	0	* *
Alleghany	44	* *	46	* *	14	* *	0	* *	0	* *
Anson	190	34.2 (61)	210	20.5 (55)	59	* *	0	* *	0	* *
Ashe	145	53.1 (81)	153	37.3 (82)	70	* *	0	* *	0	* *
Avery	105	25.7 (37)	30	* *	4	* *	0	* *	0	* *
Beaufort	298	30.2 (52)	282	17.7 (40)	59	* *	0	* *	0	* *
Bertie	147	18.4 (22)	118	7.6 (3)	18	* *	0	* *	0	* *
Bladen	289	18.0 (20)	172	17.4 (38)	22	* *	0	* *	0	* *
Brunswick	417	32.9 (56)	440	24.5 (67)	96	* *	0	* *	0	* *
Buncombe	1,231	27.3 (42)	1,134	20.3 (54)	265	22.3 (28)	0	* *	0	* *
Burke	325	53.8 (82)	587	29.0 (79)	162	21.0 (29)	0	* *	0	* *
Cabarrus	1,034	23.9 (32)	1,186	13.7 (26)	222	38.7 (8)	0	* *	0	* *
Caldwell	421	34.7 (64)	499	27.5 (76)	127	19.7 (31)	0	* *	0	* *
Camden	25	* *	26	* *	7	* *	0	* *	0	* *

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one county's rate to other counties. Rank 1 = best rate.

(4) Among women who smoked 3 months prior to pregnancy.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

Table 7B
2007 Pregnancy Nutrition Surveillance (1)
North Carolina
Comparison of Smoking Indicators by County

Comparison of Smoking Indicators (2, 3)										
County	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)		Smoking in Household			
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy				Prenatal		Postpartum	
	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Carteret	260	48.1 (80)	312	23.4 (62)	107	44.9 (5)	0	* *	0	* *
Caswell	98	* *	105	24.8 (68)	20	* *	0	* *	0	* *
Catawba	936	32.9 (56)	869	18.3 (45)	233	35.2 (11)	0	* *	0	* *
Chatham	283	12.7 (6)	304	14.1 (27)	25	* *	0	* *	0	* *
Cherokee	121	57.9 (85)	119	47.1 (83)	61	* *	0	* *	0	* *
Chowan	105	35.2 (66)	104	19.2 (50)	34	* *	0	* *	0	* *
Clay	28	* *	26	* *	11	* *	0	* *	0	* *
Cleveland	571	44.0 (77)	581	22.0 (60)	182	30.2 (16)	0	* *	0	* *
Columbus	366	26.5 (39)	228	26.3 (75)	46	* *	0	* *	0	* *
Craven	671	29.2 (48)	756	13.6 (24)	137	36.5 (9)	0	* *	0	* *
Cumberland	2,347	31.9 (55)	2,985	11.9 (12)	589	47.4 (3)	0	* *	0	* *
Currituck	69	* *	74	* *	37	* *	0	* *	0	* *
Dare	121	28.1 (45)	157	18.5 (47)	26	* *	0	* *	0	* *
Davidson	853	40.1 (74)	944	24.8 (68)	301	30.2 (16)	0	* *	0	* *
Davie	168	26.2 (38)	192	19.3 (51)	39	* *	0	* *	0	* *

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one county's rate to other counties. Rank 1 = best rate.

(4) Among women who smoked 3 months prior to pregnancy.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

Table 7B
2007 Pregnancy Nutrition Surveillance (1)
North Carolina
Comparison of Smoking Indicators by County

Comparison of Smoking Indicators (2, 3)										
County	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)		Smoking in Household			
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy				Prenatal		Postpartum	
	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Duplin	469	10.9 (4)	573	8.7 (5)	50	* *	0	* *	0	* *
Durham	1,269	15.3 (13)	1,746	8.2 (4)	169	46.2 (4)	0	* *	0	* *
Edgecombe	286	14.7 (10)	572	13.3 (20)	38	* *	0	* *	0	* *
Forsyth	1,713	15.6 (15)	2,123	10.7 (9)	213	33.8 (12)	0	* *	0	* *
Franklin	255	29.4 (49)	273	13.2 (19)	55	* *	0	* *	0	* *
Gaston	1,220	33.6 (59)	1,328	19.4 (52)	342	33.0 (13)	0	* *	0	* *
Gates	22	* *	29	* *	4	* *	0	* *	0	* *
Graham	64	* *	63	* *	19	* *	0	* *	0	* *
Granville	265	29.1 (47)	296	16.9 (36)	65	* *	0	* *	0	* *
Greene	184	17.9 (18)	212	12.3 (16)	29	* *	0	* *	0	* *
Guilford	2,219	15.3 (13)	2,486	12.0 (13)	266	30.1 (18)	0	* *	0	* *
Halifax	459	31.4 (54)	523	16.6 (35)	139	51.8 (2)	0	* *	0	* *
Harnett	680	28.7 (46)	675	17.8 (42)	148	24.3 (27)	0	* *	0	* *
Haywood	266	56.8 (84)	256	36.3 (81)	121	27.3 (24)	0	* *	0	* *
Henderson	296	17.6 (17)	345	13.6 (24)	26	* *	0	* *	0	* *

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one county's rate to other counties. Rank 1 = best rate.

(4) Among women who smoked 3 months prior to pregnancy.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

Table 7B
2007 Pregnancy Nutrition Surveillance (1)
North Carolina
Comparison of Smoking Indicators by County

Comparison of Smoking Indicators (2, 3)										
County	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)	Smoking in Household				
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy			Prenatal		Postpartum		
	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Hertford	180	23.9 (32)	147	12.2 (14)	32	* *	0	* *	0	* *
Hoke	333	21.6 (26)	217	8.8 (6)	35	* *	0	* *	0	* *
Hyde	25	* *	25	* *	6	* *	0	* *	0	* *
Iredell	806	30.1 (51)	916	21.1 (56)	215	20.9 (30)	0	* *	0	* *
Jackson	165	34.5 (62)	225	25.3 (72)	50	* *	0	* *	0	* *
Johnston	640	20.5 (23)	886	12.8 (17)	98	* *	0	* *	0	* *
Jones	70	* *	12	* *	2	* *	0	* *	0	* *
Lee	422	24.9 (36)	454	13.4 (22)	85	* *	0	* *	0	* *
Lenoir	382	8.4 (2)	393	17.6 (39)	23	* *	0	* *	0	* *
Lincoln	351	39.9 (73)	318	25.2 (71)	102	36.3 (10)	0	* *	0	* *
Macon	255	34.9 (65)	296	23.6 (64)	77	* *	0	* *	0	* *
Madison	215	43.7 (76)	138	27.5 (76)	45	* *	0	* *	0	* *
Martin	100	39.0 (72)	37	* *	6	* *	0	* *	0	* *
Mcdowell	171	23.4 (30)	178	16.9 (36)	34	* *	0	* *	0	* *
Mecklenbur	2,575	4.8 (1)	3,641	6.8 (2)	96	* *	0	* *	0	* *

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one county's rate to other counties. Rank 1 = best rate.
(4) Among women who smoked 3 months prior to pregnancy.
* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 7B
2007 Pregnancy Nutrition Surveillance (1)
North Carolina
Comparison of Smoking Indicators by County

Comparison of Smoking Indicators (2, 3)										
County	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)	Smoking in Household				
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy			Prenatal		Postpartum		
	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Mitchell	78	* *	76	* *	32	* *	0	* *	0	* *
Montgomery	197	27.9 (44)	112	21.4 (58)	25	* *	0	* *	0	* *
Moore	156	14.1 (8)	195	9.2 (7)	7	* *	0	* *	0	* *
Nash	608	21.2 (25)	668	15.7 (32)	117	28.2 (21)	0	* *	0	* *
New Hanove	531	21.1 (24)	957	15.6 (31)	82	* *	0	* *	0	* *
Northhampt	131	22.9 (28)	121	14.9 (28)	22	* *	0	* *	0	* *
Onslow	1,714	34.6 (63)	1,782	12.8 (17)	461	57.0 (1)	0	* *	0	* *
Orange	387	15.0 (11)	396	16.4 (34)	38	* *	0	* *	0	* *
Pamlico	65	* *	16	* *	5	* *	0	* *	0	* *
Pasquotank	189	11.6 (5)	263	9.5 (8)	19	* *	0	* *	0	* *
Pender	268	24.6 (35)	332	18.4 (46)	62	* *	0	* *	0	* *
Perquimans	54	* *	64	* *	9	* *	0	* *	0	* *
Person	230	42.2 (75)	166	25.9 (74)	55	* *	0	* *	0	* *
Pitt	985	22.3 (27)	1,033	12.2 (14)	173	39.9 (6)	0	* *	0	* *
Polk	41	* *	28	* *	4	* *	0	* *	0	* *

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one county's rate to other counties. Rank 1 = best rate.

(4) Among women who smoked 3 months prior to pregnancy.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

Table 7B
2007 Pregnancy Nutrition Surveillance (1)
North Carolina
Comparison of Smoking Indicators by County

Comparison of Smoking Indicators (2, 3)										
County	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)		Smoking in Household			
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy				Prenatal		Postpartum	
	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Randolph	792	29.9 (50)	876	15.8 (33)	189	25.4 (25)	0	* *	0	* *
Richmond	428	38.3 (71)	457	23.0 (61)	136	39.0 (7)	0	* *	0	* *
Robeson	1,201	17.2 (16)	1,152	18.7 (48)	143	8.4 (32)	0	* *	0	* *
Rockingham	580	36.9 (70)	613	24.3 (66)	190	28.9 (20)	0	* *	0	* *
Rowan	597	33.8 (60)	883	19.4 (52)	175	25.1 (26)	0	* *	0	* *
Rutherford	196	35.2 (66)	285	25.6 (73)	44	* *	0	* *	0	* *
Sampson	471	14.2 (9)	544	11.0 (11)	56	* *	0	* *	0	* *
Scotland	398	23.1 (29)	357	21.3 (57)	68	* *	0	* *	0	* *
Stanly	343	26.5 (39)	314	23.6 (64)	61	* *	0	* *	0	* *
Stokes	175	56.6 (83)	214	34.6 (80)	82	* *	0	* *	0	* *
Surry	405	27.4 (43)	411	23.4 (62)	80	* *	0	* *	0	* *
Swain	118	69.5 (86)	81	* *	30	* *	0	* *	0	* *
Transylvan	143	46.9 (79)	91	* *	31	* *	0	* *	0	* *
Tyrrell	40	* *	33	* *	2	* *	0	* *	0	* *
Union	423	13.2 (7)	782	13.4 (22)	26	* *	0	* *	0	* *

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one county's rate to other counties. Rank 1 = best rate.
(4) Among women who smoked 3 months prior to pregnancy.
* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 7B
2007 Pregnancy Nutrition Surveillance (1)
North Carolina
Comparison of Smoking Indicators by County

Comparison of Smoking Indicators (2, 3)										
County	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)		Smoking in Household			
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy				Prenatal		Postpartum	
	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Vance	414	15.0 (11)	407	18.7 (48)	51	* *	0	* *	0	* *
Wake	2,151	9.9 (3)	4,016	5.8 (1)	180	29.4 (19)	0	* *	0	* *
Warren	145	24.1 (34)	131	15.3 (30)	26	* *	0	* *	0	* *
Washington	134	17.9 (18)	121	10.7 (9)	20	* *	0	* *	0	* *
Watauga	121	44.6 (78)	101	17.8 (42)	22	* *	0	* *	0	* *
Wayne Coun	950	23.4 (30)	685	13.3 (20)	119	31.9 (14)	0	* *	0	* *
Wilkes	451	36.1 (69)	399	24.8 (68)	123	27.6 (23)	0	* *	0	* *
Wilson Cou	662	18.0 (20)	621	15.1 (29)	100	28.0 (22)	0	* *	0	* *
Yadkin	222	31.1 (53)	249	18.1 (44)	60	* *	0	* *	0	* *
Yancey	83	* *	49	* *	13	* *	0	* *	0	* *
Unknown Counties	232	26.7 (41)	215	17.7 (40)	51	* *	0	* *	0	* *
North Carolina	46,721	25.2	53,527	15.4	9,076	32.1	0	*	0	*
Nation (Prior Year)	847,411	27.2	692,984	16.8	156,923	35.0	514,582	22.5	443,951	24.1

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one county's rate to other counties. Rank 1 = best rate.
(4) Among women who smoked 3 months prior to pregnancy.
* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 8B
2007 Pregnancy Nutrition Surveillance ⁽¹⁾
North Carolina
Comparison of Infant Health Indicators by County

Comparison of Infant Health Indicators ^(2, 3)										
County	Birthweight				Preterm ⁽⁶⁾		Full Term LBW ⁽⁷⁾		Ever Breastfed	
	Low ⁽⁴⁾		High ⁽⁵⁾		Number	% ^(Rank)	Number	% ^(Rank)	Number	% ^(Rank)
	Number	% ^(Rank)	% ^(Rank)							
Alamance	1,020	7.5 (16)	9.3 (78)		1,015	12.1 (20)	892	2.7 (4)	1,016	55.7 (45)
Alexander	201	9.5 (43)	6.5 (49)		199	13.1 (31)	173	4.6 (50)	100	94.0 (2)
Alleghany	46	* * * *			46	* *	42	* *	45	* *
Anson	210	10.0 (51)	5.2 (20)		208	16.3 (62)	174	4.0 (36)	209	34.4 (74)
Ashe	154	6.5 (6)	4.5 (12)		153	9.2 (3)	139	3.6 (21)	151	55.6 (46)
Avery	31	* * * *			28	* *	22	* *	31	* *
Beaufort	287	9.8 (49)	5.6 (23)		281	12.8 (25)	245	6.5 (73)	283	47.0 (60)
Bertie	115	11.3 (70)	5.2 (20)		113	15.0 (52)	96	* *	116	15.5 (81)
Bladen	171	9.4 (42)	7.6 (66)		168	17.9 (74)	138	5.8 (69)	171	52.6 (54)
Brunswick	440	8.6 (32)	7.0 (58)		436	11.0 (10)	388	5.4 (63)	431	39.2 (69)
Buncombe	1,135	9.6 (45)	6.5 (49)		1,118	11.3 (15)	992	4.5 (46)	1,108	81.1 (5)
Burke	587	8.3 (27)	5.8 (31)		575	14.8 (47)	490	5.7 (67)	539	63.3 (25)
Cabarrus	1,196	10.1 (53)	5.7 (26)		1,176	15.4 (56)	995	3.4 (17)	1,194	70.1 (14)
Caldwell	499	8.6 (32)	6.2 (43)		489	12.3 (21)	429	5.4 (63)	498	55.2 (50)

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one county's rate to other counties. Rank 1 = best rate.
(4) <2500 g.
(5) >4000 g.
(6) <37 weeks gestation.
(7) 37 or more weeks gestation and <2500 g.
* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 8B
2007 Pregnancy Nutrition Surveillance ⁽¹⁾
North Carolina
Comparison of Infant Health Indicators by County

Comparison of Infant Health Indicators ^(2, 3)										
County	Birthweight				Preterm ⁽⁶⁾		Full Term LBW ⁽⁷⁾		Ever Breastfed	
	Low ⁽⁴⁾		High ⁽⁵⁾		Number	%	Number	%	Number	%
	Number	% (Rank)	% (Rank)	Number						
Camden	26	* * * *			26	* *	22	* *	23	* *
Carteret	316	7.0 (9)	7.9 (69)		316	7.9 (2)	290	4.8 (53)	313	64.5 (24)
Caswell	105	12.4 (76)	5.7 (26)		105	9.5 (4)	95	* *	105	67.6 (19)
Catawba	871	8.6 (32)	6.4 (47)		859	12.3 (21)	753	4.5 (46)	868	68.9 (18)
Chatham	309	7.8 (18)	9.4 (80)		301	12.3 (21)	264	3.0 (11)	306	78.8 (7)
Cherokee	120	5.8 (3)	4.2 (9)		119	13.4 (32)	103	1.9 (1)	119	56.3 (41)
Chowan	105	10.5 (62)	6.7 (54)		104	15.4 (56)	88	* *	104	26.0 (79)
Clay	26	* * * *			26	* *	25	* *	26	* *
Cleveland	588	7.1 (11)	9.2 (77)		583	13.7 (36)	503	3.0 (11)	580	29.5 (78)
Columbus	230	12.2 (73)	7.8 (68)		223	17.0 (69)	184	3.8 (27)	231	42.9 (66)
Craven	764	8.0 (24)	8.9 (74)		754	10.7 (8)	673	4.2 (42)	746	60.5 (31)
Cumberland	2,982	10.5 (62)	6.6 (52)		2,961	16.4 (65)	2,475	4.6 (50)	2,965	62.1 (29)
Currituck	76	* * * *			75	* *	68	* *	69	* *
Dare	162	5.6 (1)	13.6 (83)		161	6.8 (1)	150	4.0 (36)	128	98.4 (1)

- (1) Reporting period is January 1 through December 31.
- (2) Excludes records with unknown data or errors.
- (3) Rank compares one county's rate to other counties. Rank 1 = best rate.
- (4) <2500 g.
- (5) >4000 g.
- (6) <37 weeks gestation.
- (7) 37 or more weeks gestation and <2500 g.
- * Percentages are not calculated if <100 records are available for analysis after exclusions.

Table 8B
2007 Pregnancy Nutrition Surveillance ⁽¹⁾
North Carolina
Comparison of Infant Health Indicators by County

Comparison of Infant Health Indicators ^(2, 3)										
County	Birthweight				Preterm (6)		Full Term LBW (7)		Ever Breastfed	
	Low (4)		High (5)							
	Number	% (Rank)	% (Rank)		Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Davidson	946	9.5 (43)	5.9 (35)		924	13.5 (34)	799	3.9 (32)	933	57.1 (37)
Davie	192	7.8 (18)	7.3 (63)		190	18.9 (79)	154	3.9 (32)	186	55.4 (48)
Duplin	571	7.0 (9)	7.9 (69)		569	10.7 (8)	508	3.1 (13)	563	51.5 (56)
Durham	1,771	10.4 (60)	7.1 (61)		1,745	14.8 (47)	1,485	5.5 (65)	1,709	69.9 (15)
Edgecombe	569	9.8 (49)	4.6 (13)		558	15.8 (58)	470	4.0 (36)	561	30.5 (77)
Forsyth	2,127	10.7 (65)	6.1 (40)		2,094	17.2 (71)	1,733	4.4 (44)	2,110	76.7 (10)
Franklin	280	11.4 (71)	8.9 (74)		276	13.0 (30)	240	7.5 (76)	280	49.6 (59)
Gaston	1,347	10.8 (67)	5.8 (31)		1,306	14.0 (39)	1,123	5.3 (62)	1,346	45.5 (64)
Gates	30	* * *	* *		28	* *	27	* *	30	* *
Graham	64	* * *	* *		64	* *	59	* *	27	* *
Granville	298	10.7 (65)	4.4 (10)		291	14.8 (47)	248	4.8 (53)	285	65.3 (22)
Greene	214	7.9 (20)	6.1 (40)		211	16.1 (60)	177	2.8 (7)	215	56.3 (41)
Guilford	2,489	10.3 (59)	5.0 (18)		2,443	14.4 (43)	2,092	4.5 (46)	2,444	65.6 (21)
Halifax	523	13.8 (82)	4.4 (10)		516	22.7 (82)	399	5.0 (57)	515	34.6 (73)

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one county's rate to other counties. Rank 1 = best rate.
(4) <2500 g.
(5) >4000 g.
(6) <37 weeks gestation.
(7) 37 or more weeks gestation and <2500 g.
* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 8B
2007 Pregnancy Nutrition Surveillance ⁽¹⁾
North Carolina
Comparison of Infant Health Indicators by County

Comparison of Infant Health Indicators ^(2, 3)									
County	Birthweight			Preterm ⁽⁶⁾		Full Term LBW ⁽⁷⁾		Ever Breastfed	
	Low ⁽⁴⁾	High ⁽⁵⁾							
	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Harnett	676	8.4 (29)	6.2 (43)	668	15.3 (54)	566	3.9 (32)	665	56.4 (38)
Haywood	256	12.5 (79)	5.5 (22)	254	16.5 (67)	212	3.8 (27)	251	70.5 (13)
Henderson	348	7.2 (12)	7.5 (65)	344	12.8 (25)	300	3.7 (26)	346	84.4 (4)
Hertford	147	11.6 (72)	6.1 (40)	142	17.6 (72)	117	3.4 (17)	147	35.4 (72)
Hoke	217	10.1 (53)	6.0 (37)	213	17.8 (73)	175	4.6 (50)	217	52.5 (55)
Hyde	25	* * * *		25	* *	23	* *	25	* *
Iredell	918	10.6 (64)	7.0 (58)	913	15.8 (58)	769	4.3 (43)	907	55.6 (46)
Jackson	225	6.2 (5)	11.6 (81)	225	11.1 (13)	200	2.5 (3)	219	69.4 (17)
Johnston	887	8.2 (26)	6.2 (43)	885	15.3 (54)	750	3.2 (14)	879	52.9 (53)
Jones	12	* * * *		11	* *	9	* *	12	* *
Lee	455	8.8 (37)	4.8 (15)	447	14.8 (47)	381	3.9 (32)	456	59.4 (34)
Lenoir	400	7.3 (14)	5.8 (31)	382	12.3 (21)	335	3.6 (21)	396	45.5 (64)
Lincoln	318	10.1 (53)	7.2 (62)	312	12.8 (25)	271	5.2 (60)	319	56.4 (38)
Macon	296	8.4 (29)	3.7 (7)	291	11.0 (10)	259	5.0 (57)	292	58.9 (35)

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one county's rate to other counties. Rank 1 = best rate.
(4) <2500 g.
(5) >4000 g.
(6) <37 weeks gestation.
(7) 37 or more weeks gestation and <2500 g.
* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 8B
2007 Pregnancy Nutrition Surveillance ⁽¹⁾
North Carolina
Comparison of Infant Health Indicators by County

Comparison of Infant Health Indicators ^(2, 3)										
County	Birthweight				Preterm ⁽⁶⁾		Full Term LBW ⁽⁷⁾		Ever Breastfed	
	Low ⁽⁴⁾		High ⁽⁵⁾		Number	%	Number	%	Number	%
	Number	% (Rank)	% (Rank)	(Rank)						
Madison	139	7.9 (20)	3.6 (6)		136	11.0 (10)	121	3.3 (16)	138	74.6 (12)
Martin	37	* * *	* *		36	* *	31	* *	29	* *
Mcdowell	179	12.3 (74)	3.4 (5)		177	16.4 (65)	148	7.4 (75)	175	37.7 (70)
Mecklenbur	3,646	10.4 (60)	5.8 (31)		3,615	14.5 (45)	3,091	3.8 (27)	3,593	65.8 (20)
Mitchell	75	* * *	* *		71	* *	67	* *	74	* *
Montgomery	112	8.0 (24)	2.7 (3)		106	14.2 (42)	91	* *	112	55.4 (48)
Moore	195	9.7 (46)	5.6 (23)		189	10.1 (6)	170	5.9 (70)	194	59.8 (32)
Nash	671	8.6 (32)	7.0 (58)		652	11.5 (17)	577	3.8 (27)	666	50.0 (57)
New Hanove	954	10.2 (57)	7.4 (64)		950	14.0 (39)	817	4.4 (44)	950	55.9 (44)
Northhampt	121	12.4 (76)	3.3 (4)		118	16.9 (68)	98	* *	116	17.2 (80)
Onslow	1,779	6.9 (8)	9.3 (78)		1,765	10.6 (7)	1,578	2.7 (4)	1,702	65.3 (22)
Orange	403	6.7 (7)	6.5 (49)		389	11.1 (13)	346	3.5 (19)	397	79.1 (6)
Pamlico	18	* * *	* *		17	* *	14	* *	17	* *
Pasquotank	265	8.7 (36)	5.7 (26)		264	16.3 (62)	221	4.1 (40)	251	7.6 (82)

- (1) Reporting period is January 1 through December 31.
- (2) Excludes records with unknown data or errors.
- (3) Rank compares one county's rate to other counties. Rank 1 = best rate.
- (4) <2500 g.
- (5) >4000 g.
- (6) <37 weeks gestation.
- (7) 37 or more weeks gestation and <2500 g.
- * Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 8B
2007 Pregnancy Nutrition Surveillance ⁽¹⁾
North Carolina
Comparison of Infant Health Indicators by County

Comparison of Infant Health Indicators ^(2, 3)										
County	Birthweight				Preterm ⁽⁶⁾		Full Term LBW ⁽⁷⁾		Ever Breastfed	
	Low ⁽⁴⁾		High ⁽⁵⁾		Number	%	Number	%	Number	%
	Number	% (Rank)	Number	% (Rank)						
Pender	331	10.0 (51)	6.3 (46)		329	11.9 (18)	290	4.8 (53)	327	56.3 (41)
Perquimans	65	* * *	* *		64	* *	56	* *	65	* *
Person	172	9.3 (40)	12.2 (82)		167	18.6 (78)	136	2.9 (8)	170	54.1 (51)
Pitt	1,059	13.5 (81)	5.9 (35)		1,032	18.5 (77)	840	5.0 (57)	1,046	53.0 (52)
Polk	28	* * *	* *		28	* *	26	* *	28	* *
Randolph	881	7.9 (20)	8.3 (72)		874	15.0 (52)	743	2.7 (4)	877	69.7 (16)
Richmond	458	7.9 (20)	4.6 (13)		452	13.9 (37)	389	4.1 (40)	455	33.4 (75)
Robeson	1,160	10.9 (69)	5.7 (26)		1,122	17.1 (70)	930	5.7 (67)	1,149	36.6 (71)
Rockingham	607	9.7 (46)	6.4 (47)		609	12.8 (25)	531	4.5 (46)	607	49.9 (58)
Rowan	887	10.8 (67)	6.0 (37)		869	16.3 (62)	727	4.0 (36)	887	59.6 (33)
Rutherford	288	8.3 (27)	6.9 (56)		283	11.3 (15)	251	3.6 (21)	287	61.0 (30)
Sampson	544	7.2 (12)	5.7 (26)		540	13.9 (37)	465	3.2 (14)	544	40.3 (67)
Scotland	358	12.3 (74)	5.6 (23)		350	13.4 (32)	303	5.9 (70)	356	30.9 (76)
Stanly	314	10.2 (57)	4.1 (8)		311	14.5 (45)	266	4.9 (56)	305	76.4 (11)

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one county's rate to other counties. Rank 1 = best rate.
(4) <2500 g.
(5) >4000 g.
(6) <37 weeks gestation.
(7) 37 or more weeks gestation and <2500 g.
* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 8B
2007 Pregnancy Nutrition Surveillance ⁽¹⁾
North Carolina
Comparison of Infant Health Indicators by County

Comparison of Infant Health Indicators ^(2, 3)										
County	Birthweight				Preterm ⁽⁶⁾		Full Term LBW ⁽⁷⁾		Ever Breastfed	
	Low ⁽⁴⁾		High ⁽⁵⁾		Number	%	Number	%	Number	%
	Number	% (Rank)	% (Rank)	(Rank)						
Stokes	213	12.7 (80)	8.9 (74)		210	14.8 (47)	179	7.3 (74)	211	56.4 (38)
Surry	411	7.3 (14)	8.8 (73)		407	12.0 (19)	358	3.6 (21)	378	39.9 (68)
Swain	80	*	*	*	78	*	70	*	73	*
Transylvan	91	*	*	*	90	*	82	*	90	*
Tyrrell	36	*	*	*	35	*	29	*	35	*
Union	784	5.7 (2)	6.8 (55)		779	10.0 (5)	701	1.9 (1)	727	76.9 (9)
Vance	411	12.4 (76)	5.1 (19)		402	20.6 (80)	319	5.6 (66)	318	62.3 (28)
Wake	4,025	7.7 (17)	7.7 (67)		3,959	14.4 (43)	3,388	2.9 (8)	3,501	90.4 (3)
Warren	133	16.5 (83)	2.3 (1)		130	22.3 (81)	101	5.9 (70)	129	46.5 (62)
Washington	122	9.0 (38)	6.6 (52)		120	18.3 (75)	98	*	122	45.9 (63)
Watauga	101	5.9 (4)	6.9 (56)		98	*	89	*	98	*
Wayne Coun	691	9.3 (40)	8.2 (71)		680	12.9 (29)	591	3.6 (21)	661	58.4 (36)
Wilkes	399	9.0 (38)	4.8 (15)		398	14.1 (41)	342	3.5 (19)	391	62.9 (27)
Wilson Cou	620	9.7 (46)	4.8 (15)		608	18.4 (76)	496	5.2 (60)	617	47.0 (60)

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one county's rate to other counties. Rank 1 = best rate.
(4) <2500 g.
(5) >4000 g.
(6) <37 weeks gestation.
(7) 37 or more weeks gestation and <2500 g.
* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 8B
2007 Pregnancy Nutrition Surveillance ⁽¹⁾
North Carolina
Comparison of Infant Health Indicators by County

Comparison of Infant Health Indicators ^(2, 3)										
County	Birthweight				Preterm (6)		Full Term LBW (7)		Ever Breastfed	
	Low (4)		High (5)							
	Number	% (Rank)	%	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Yadkin	249	8.4 (29)	6.0	(37)	249	16.1 (60)	209	2.9 (8)	248	78.6 (8)
Yancey	49	* * *	*	*	48	* *	40	* *	48	* *
Unknown Counties	217	10.1 (53)	2.3	(1)	215	13.5 (34)	186	3.8 (27)	206	63.1 (26)
North Carolina	53,751	9.4	6.6		52,966	14.3	45,390	4.1	52,180	61.1
Nation (Prior Year)	972,241	8.3	6.8		873,261	12.1	802,123	3.7	897,379	64.2

- (1) Reporting period is January 1 through December 31.
- (2) Excludes records with unknown data or errors.
- (3) Rank compares one county's rate to other counties. Rank 1 = best rate.
- (4) <2500 g.
- (5) >4000 g.
- (6) <37 weeks gestation.
- (7) 37 or more weeks gestation and <2500 g.
- * Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 3B
2005 - 2007 Pregnancy Nutrition Surveillance ⁽¹⁾
North Carolina
Comparison of Racial and Ethnic Distribution by County
Combined 3 Years

Comparison of Racial and Ethnic Distribution ⁽²⁾								
County	Number Included in Analysis	White, Not Hispanic %	Black, Not Hispanic %	Hispanic %	American Indian/Alaskan Native %	Asian/Pacific Islander %	Multiple Races %	All Other/Unknown %
Alamance	3,243	39.2	25.3	33.9	0.2	1.2	0.0	0.2
Alexander	719	80.8	6.5	9.3	0.4	2.8	0.0	0.1
Alleghany	152	81.6	1.3	16.4	0.0	0.7	0.0	0.0
Anson	702	29.2	67.2	2.4	0.1	1.0	0.0	0.0
Ashe	496	92.3	0.6	6.7	0.2	0.2	0.0	0.0
Avery	312	86.9	0.0	12.5	0.0	0.6	0.0	0.0
Beaufort	1,119	37.7	42.8	19.1	0.1	0.3	0.0	0.0
Bertie	530	16.4	82.3	0.9	0.2	0.2	0.0	0.0
Bladen	915	37.2	42.5	16.9	3.1	0.3	0.0	0.0
Brunswick	1,782	65.8	18.2	13.9	1.2	0.6	0.0	0.3
Buncombe	4,205	67.7	13.4	16.9	0.2	1.7	0.0	0.0
Burke	1,741	65.6	5.9	18.6	0.2	9.6	0.0	0.1
Cabarrus	3,463	41.4	21.5	35.6	0.1	1.2	0.0	0.0
Caldwell	1,647	82.2	6.9	9.7	0.3	0.5	0.0	0.4
Camden	103	67.0	27.2	1.9	0.0	3.9	0.0	0.0
Carteret	1,045	76.6	11.5	10.2	1.2	0.5	0.0	0.0
Caswell	374	51.1	40.1	7.8	0.5	0.0	0.0	0.5
Catawba	3,031	55.7	13.5	22.2	0.4	8.1	0.0	0.1
Chatham	1,007	29.5	15.7	54.0	0.2	0.3	0.0	0.3
Cherokee	532	90.2	2.6	4.1	2.3	0.4	0.0	0.4

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 3B
2005 - 2007 Pregnancy Nutrition Surveillance ⁽¹⁾
North Carolina
Comparison of Racial and Ethnic Distribution by County
Combined 3 Years

Comparison of Racial and Ethnic Distribution ⁽²⁾								
County	Number Included in Analysis	White, Not Hispanic %	Black, Not Hispanic %	Hispanic %	American Indian/Alaskan Native %	Asian/Pacific Islander %	Multiple Races %	All Other/Unknown %
Chowan	355	36.9	58.0	5.1	0.0	0.0	0.0	0.0
Clay	134	89.6	1.5	7.5	1.5	0.0	0.0	0.0
Cleveland	2,159	57.9	36.7	4.0	0.2	0.7	0.0	0.4
Columbus	1,347	44.5	42.8	9.9	2.6	0.1	0.0	0.1
Craven	2,683	53.3	33.5	10.0	0.4	2.6	0.0	0.1
Cumberland	9,839	39.3	43.2	12.8	2.1	2.6	0.0	0.0
Currituck	225	78.7	12.9	8.0	0.0	0.4	0.0	0.0
Dare	550	67.1	3.8	27.8	0.2	0.9	0.0	0.2
Davidson	2,973	65.7	13.6	18.0	0.3	2.2	0.0	0.2
Davie	621	67.6	7.7	24.3	0.2	0.0	0.0	0.2
Duplin	1,712	30.4	26.9	42.1	0.5	0.2	0.0	0.0
Durham	5,434	8.4	50.9	38.6	0.3	1.6	0.0	0.2
Edgecombe	1,692	21.4	72.9	5.5	0.1	0.1	0.0	0.0
Forsyth	7,146	22.9	38.8	37.3	0.2	0.8	0.0	0.1
Franklin	1,014	41.1	37.4	20.8	0.4	0.1	0.0	0.2
Gaston	4,375	57.9	25.2	15.2	0.1	1.3	0.0	0.3
Gates	93	*	*	*	*	*	*	*
Graham	229	82.5	0.0	4.4	12.7	0.0	0.0	0.4
Granville	1,004	37.0	46.5	15.5	0.3	0.7	0.0	0.0
Greene	591	25.0	48.7	25.9	0.2	0.2	0.0	0.0

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

Table 3B
2005 - 2007 Pregnancy Nutrition Surveillance ⁽¹⁾
North Carolina
Comparison of Racial and Ethnic Distribution by County
Combined 3 Years

Comparison of Racial and Ethnic Distribution ⁽²⁾								
County	Number Included in Analysis	White, Not Hispanic	Black, Not Hispanic	Hispanic	American Indian/Alaskan Native	Asian/Pacific Islander	Multiple Races	All Other/Unknown
		%	%	%	%	%	%	%
Guilford	8,723	22.2	54.2	18.5	0.7	4.3	0.0	0.1
Halifax	1,609	22.6	67.5	2.9	6.8	0.1	0.0	0.1
Harnett	2,395	46.6	29.8	21.3	1.5	0.5	0.0	0.2
Haywood	933	86.3	2.6	9.9	1.0	0.3	0.0	0.0
Henderson	1,498	52.7	5.0	40.4	0.1	1.7	0.0	0.0
Hertford	657	16.6	81.4	1.4	0.2	0.5	0.0	0.0
Hoke	1,313	27.6	34.4	22.9	13.9	1.1	0.0	0.0
Hyde	107	54.2	32.7	13.1	0.0	0.0	0.0	0.0
Iredell	2,888	53.3	25.6	18.7	0.1	2.3	0.0	0.0
Jackson	788	62.1	3.0	11.8	21.1	1.9	0.0	0.1
Johnston	3,028	39.7	22.8	36.6	0.6	0.4	0.0	0.0
Jones	232	52.6	34.9	11.6	0.4	0.0	0.0	0.4
Lee	1,543	31.0	26.8	41.3	0.3	0.5	0.0	0.1
Lenoir	1,481	30.9	54.8	13.6	0.1	0.4	0.0	0.1
Lincoln	1,294	77.4	8.7	13.0	0.2	0.6	0.0	0.0
Macon	1,013	83.5	2.8	12.3	0.4	0.9	0.0	0.1
Madison	692	76.0	0.9	20.4	1.4	1.0	0.0	0.3
Martin	378	93.1	0.8	5.0	0.5	0.5	0.0	0.0
Mcdowell	619	30.2	61.4	7.4	0.0	1.0	0.0	0.0
Mecklenbur	12,757	13.3	53.5	29.7	0.3	2.9	0.0	0.2

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 3B
2005 - 2007 Pregnancy Nutrition Surveillance ⁽¹⁾
North Carolina
Comparison of Racial and Ethnic Distribution by County
Combined 3 Years

Comparison of Racial and Ethnic Distribution ⁽²⁾								
County	Number Included in Analysis	White, Not Hispanic	Black, Not Hispanic	Hispanic	American Indian/Alaskan Native	Asian/Pacific Islander	Multiple Races	All Other/Unknown
		%	%	%	%	%	%	%
Mitchell	261	89.3	0.0	10.7	0.0	0.0	0.0	0.0
Montgomery	743	39.2	22.5	35.3	0.3	2.8	0.0	0.0
Moore	1,240	51.1	27.5	18.3	2.0	0.9	0.0	0.2
Nash	2,107	30.3	53.4	15.4	0.5	0.3	0.0	0.1
New Hanove	3,277	42.8	36.6	18.8	0.4	0.8	0.0	0.5
Northhampt	544	17.5	79.4	2.6	0.4	0.2	0.0	0.0
Onslow	6,273	65.6	18.8	11.6	0.9	2.8	0.0	0.2
Orange	1,391	30.5	29.0	36.7	0.6	2.7	0.0	0.4
Pamlico	237	66.2	24.5	8.0	0.4	0.8	0.0	0.0
Pasquotank	905	36.0	53.0	9.7	0.2	0.9	0.0	0.1
Pender	930	51.1	26.2	22.0	0.2	0.3	0.0	0.1
Perquimans	204	47.1	48.0	4.4	0.0	0.5	0.0	0.0
Person	780	47.6	41.3	10.1	0.5	0.4	0.0	0.1
Pitt	3,518	27.0	60.9	11.1	0.3	0.8	0.0	0.0
Polk	198	72.2	5.6	22.2	0.0	0.0	0.0	0.0
Randolph	2,798	61.4	7.8	29.0	0.4	1.3	0.0	0.1
Richmond	1,513	46.7	36.9	12.9	2.8	0.7	0.0	0.0
Robeson	4,395	17.1	26.2	13.2	43.0	0.3	0.0	0.2
Rockingham	1,947	60.8	25.0	13.3	0.5	0.4	0.0	0.1
Rowan	2,909	51.0	25.1	22.2	0.1	1.5	0.0	0.1

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 3B
2005 - 2007 Pregnancy Nutrition Surveillance ⁽¹⁾
North Carolina
Comparison of Racial and Ethnic Distribution by County
Combined 3 Years

Comparison of Racial and Ethnic Distribution ⁽²⁾								
County	Number Included in Analysis	White, Not Hispanic	Black, Not Hispanic	Hispanic	American Indian/Alaskan Native	Asian/Pacific Islander	Multiple Races	All Other/Unknown
		%	%	%	%	%	%	%
Rutherford	1,060	73.2	16.6	9.3	0.2	0.6	0.0	0.1
Sampson	1,764	29.1	30.5	37.8	2.3	0.2	0.0	0.1
Scotland	1,265	31.0	51.9	0.5	16.1	0.4	0.0	0.1
Stanly	1,160	64.0	19.8	11.3	0.2	4.7	0.0	0.1
Stokes	741	90.6	4.7	4.0	0.3	0.4	0.0	0.0
Surry	1,548	68.0	4.6	26.6	0.3	0.6	0.0	0.0
Swain	450	57.3	0.7	3.1	37.6	1.3	0.0	0.0
Transylvan	518	84.0	6.4	8.3	0.2	1.0	0.0	0.2
Tyrrell	127	35.4	40.2	23.6	0.0	0.8	0.0	0.0
Union	2,583	41.3	27.1	30.8	0.3	0.4	0.0	0.0
Vance	1,504	26.5	58.8	13.7	0.5	0.5	0.0	0.0
Wake	12,419	19.2	40.3	37.6	0.3	2.4	0.0	0.2
Warren	511	17.2	70.8	7.8	3.3	0.4	0.0	0.4
Washington	398	25.4	69.3	5.3	0.0	0.0	0.0	0.0
Watauga	530	87.2	2.3	9.2	0.4	0.9	0.0	0.0
Wayne Coun	2,992	32.7	42.0	23.5	0.3	1.2	0.0	0.3
Wilkes	1,412	79.2	5.3	14.7	0.1	0.6	0.0	0.0
Wilson Cou	2,147	20.4	53.2	25.5	0.0	0.8	0.0	0.1
Yadkin	783	59.6	3.8	36.1	0.1	0.3	0.0	0.0
Yancey	315	89.5	0.3	9.8	0.3	0.0	0.0	0.0

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 3B
2005 - 2007 Pregnancy Nutrition Surveillance ⁽¹⁾
North Carolina
Comparison of Racial and Ethnic Distribution by County
Combined 3 Years

Comparison of Racial and Ethnic Distribution ⁽²⁾								
County	Number Included in Analysis	White, Not Hispanic %	Black, Not Hispanic %	Hispanic %	American Indian/Alaskan Native %	Asian/Pacific Islander %	Multiple Races %	All Other/Unknown %
Unknown Counties	946	39.6	31.3	25.1	2.7	1.1	0.0	0.2
North Carolina	186,585	40.7	33.8	21.7	2.0	1.7	0.0	0.1

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

Table 4B
2005 - 2007 Pregnancy Nutrition Surveillance ⁽¹⁾
North Carolina
Comparison of Age Distribution by County
Combined 3 Years

Comparison of Age Distribution ⁽²⁾								
County	Number Included in Analysis	< 15 Years %	15 - 17 Years %	18 - 19 Years %	20 - 29 Years %	30 - 39 Years %	>= 40 Years %	Unknown %
Alamance	3,243	0.7	8.3	14.9	58.8	16.5	0.9	0.0
Alexander	719	0.0	7.8	15.3	61.2	14.6	1.1	0.0
Alleghany	152	0.7	10.5	13.8	62.5	12.5	0.0	0.0
Anson	702	0.6	11.0	15.0	58.7	14.1	0.7	0.0
Ashe	496	0.2	7.7	14.1	62.1	14.7	1.2	0.0
Avery	312	0.0	4.5	12.5	58.3	23.7	1.0	0.0
Beaufort	1,119	0.8	8.4	15.4	58.5	16.4	0.5	0.0
Bertie	530	0.8	9.4	16.0	59.6	13.6	0.6	0.0
Bladen	915	0.4	9.7	13.6	58.6	16.9	0.8	0.0
Brunswick	1,782	0.4	6.7	16.0	60.1	15.4	1.3	0.0
Buncombe	4,205	0.5	7.2	14.4	59.0	17.7	1.1	0.0
Burke	1,741	0.5	10.1	15.6	57.0	16.1	0.6	0.0
Cabarrus	3,463	0.5	7.6	13.3	60.7	17.2	0.7	0.0
Caldwell	1,647	0.3	10.3	18.6	56.2	14.1	0.5	0.0
Camden	103	0.0	8.7	18.4	55.3	16.5	1.0	0.0
Carteret	1,045	0.2	6.7	14.2	63.0	14.7	1.2	0.0
Caswell	374	0.3	8.6	13.1	61.5	15.5	1.1	0.0
Catawba	3,031	0.5	8.5	16.3	57.9	15.6	1.1	0.0
Chatham	1,007	0.1	8.2	12.2	59.3	18.4	1.8	0.0

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

Table 4B
2005 - 2007 Pregnancy Nutrition Surveillance ⁽¹⁾
North Carolina
Comparison of Age Distribution by County
Combined 3 Years

Comparison of Age Distribution ⁽²⁾								
County	Number Included in Analysis	< 15 Years %	15 - 17 Years %	18 - 19 Years %	20 - 29 Years %	30 - 39 Years %	>= 40 Years %	Unknown %
Cherokee	532	0.0	4.9	13.9	63.7	15.8	1.7	0.0
Chowan	355	0.8	11.3	17.5	56.1	13.0	1.4	0.0
Clay	134	0.0	9.0	17.2	56.7	15.7	1.5	0.0
Cleveland	2,159	0.8	11.1	17.2	58.7	11.6	0.6	0.0
Columbus	1,347	0.7	9.0	17.4	57.9	13.8	1.1	0.0
Craven	2,683	0.5	6.1	15.3	66.3	11.4	0.4	0.0
Cumberland	9,839	0.4	5.3	13.0	67.4	13.2	0.7	0.0
Currituck	225	0.0	8.0	17.8	62.2	11.6	0.4	0.0
Dare	550	0.2	3.8	8.4	68.5	18.4	0.7	0.0
Davidson	2,973	0.6	8.8	16.3	59.8	13.9	0.6	0.0
Davie	621	0.6	7.7	13.8	59.4	17.2	1.1	0.0
Duplin	1,712	0.5	7.1	14.2	59.1	18.2	0.9	0.0
Durham	5,434	0.6	7.1	11.1	59.0	21.1	1.3	0.0
Edgecombe	1,692	1.0	8.7	16.3	60.5	12.6	0.9	0.0
Forsyth	7,146	0.7	8.1	12.2	60.2	17.7	1.0	0.0
Franklin	1,014	0.5	8.6	17.2	54.5	18.6	0.6	0.0
Gaston	4,375	0.7	9.3	16.0	59.0	14.2	0.7	0.0
Gates	93	*	*	*	*	*	*	*
Graham	229	0.4	10.5	9.6	65.1	13.1	1.3	0.0

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

Table 4B
2005 - 2007 Pregnancy Nutrition Surveillance ⁽¹⁾
North Carolina
Comparison of Age Distribution by County
Combined 3 Years

Comparison of Age Distribution ⁽²⁾								
County	Number Included in Analysis	< 15 Years %	15 - 17 Years %	18 - 19 Years %	20 - 29 Years %	30 - 39 Years %	>= 40 Years %	Unknown %
Granville	1,004	0.9	6.6	13.2	61.5	16.8	1.0	0.0
Greene	591	0.8	9.6	15.6	60.6	12.9	0.5	0.0
Guilford	8,723	0.5	7.0	12.4	59.3	19.6	1.2	0.0
Halifax	1,609	0.6	9.3	15.7	58.8	14.7	0.9	0.0
Harnett	2,395	0.6	8.5	14.6	58.2	17.5	0.6	0.0
Haywood	933	0.8	8.5	16.5	59.2	14.0	1.1	0.0
Henderson	1,498	0.3	7.8	12.8	61.7	16.9	0.5	0.0
Hertford	657	1.1	10.5	17.7	60.4	9.3	1.1	0.0
Hoke	1,313	1.1	7.8	11.8	62.5	15.8	1.0	0.0
Hyde	107	0.0	6.5	22.4	49.5	20.6	0.9	0.0
Iredell	2,888	0.6	8.3	15.3	59.3	15.5	1.0	0.0
Jackson	788	0.6	7.6	13.5	62.2	15.4	0.8	0.0
Johnston	3,028	0.6	7.5	13.2	58.8	18.7	1.2	0.0
Jones	232	0.4	8.2	15.1	62.5	12.9	0.9	0.0
Lee	1,543	0.5	8.0	15.1	58.3	17.0	1.2	0.0
Lenoir	1,481	0.6	10.6	16.3	56.8	14.9	0.8	0.0
Lincoln	1,294	0.5	9.4	15.2	60.0	14.6	0.4	0.0
Macon	1,013	0.3	9.2	17.1	59.6	12.9	0.9	0.0
Madison	692	0.3	8.5	13.3	61.4	15.8	0.7	0.0

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

Table 4B
2005 - 2007 Pregnancy Nutrition Surveillance ⁽¹⁾
North Carolina
Comparison of Age Distribution by County
Combined 3 Years

Comparison of Age Distribution ⁽²⁾								
County	Number Included in Analysis	< 15 Years %	15 - 17 Years %	18 - 19 Years %	20 - 29 Years %	30 - 39 Years %	>= 40 Years %	Unknown %
Martin	378	0.3	7.1	13.2	63.5	15.1	0.8	0.0
Mcdowell	619	0.6	9.9	18.9	57.0	13.4	0.2	0.0
Mecklenbur	12,757	0.5	7.5	11.9	58.9	19.9	1.3	0.0
Mitchell	261	0.0	6.5	14.2	62.8	14.9	1.5	0.0
Montgomery	743	1.1	8.3	13.7	59.5	16.8	0.5	0.0
Moore	1,240	0.4	7.7	14.0	61.6	14.6	1.6	0.0
Nash	2,107	0.7	8.3	16.3	58.2	15.4	1.0	0.0
New Hanove	3,277	0.3	6.1	13.9	62.4	16.5	0.8	0.0
Northhampt	544	0.6	10.7	17.8	54.2	15.1	1.7	0.0
Onslow	6,273	0.1	3.5	14.1	71.7	10.2	0.5	0.0
Orange	1,391	0.2	5.5	10.7	62.5	19.8	1.3	0.0
Pamlico	237	0.0	9.7	15.6	58.2	16.5	0.0	0.0
Pasquotank	905	0.1	8.3	17.6	63.5	9.9	0.6	0.0
Pender	930	0.4	6.8	14.6	60.5	16.5	1.2	0.0
Perquimans	204	1.0	8.8	19.6	55.4	13.7	1.5	0.0
Person	780	0.3	9.4	16.3	60.5	12.6	1.0	0.0
Pitt	3,518	0.6	6.8	12.4	62.6	17.0	0.6	0.0
Polk	198	0.0	8.6	11.1	62.6	16.2	1.5	0.0
Randolph	2,798	0.5	8.9	14.2	58.6	16.9	0.9	0.0

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

Table 4B
2005 - 2007 Pregnancy Nutrition Surveillance ⁽¹⁾
North Carolina
Comparison of Age Distribution by County
Combined 3 Years

Comparison of Age Distribution ⁽²⁾								
County	Number Included in Analysis	< 15 Years %	15 - 17 Years %	18 - 19 Years %	20 - 29 Years %	30 - 39 Years %	>= 40 Years %	Unknown %
Richmond	1,513	1.1	11.3	15.7	56.9	14.0	1.0	0.0
Robeson	4,395	0.8	9.1	15.3	60.2	13.9	0.7	0.0
Rockingham	1,947	0.9	8.8	17.5	58.3	14.0	0.6	0.0
Rowan	2,909	0.7	9.0	16.0	59.3	14.1	0.9	0.0
Rutherford	1,060	0.9	8.3	17.4	58.7	13.7	1.0	0.0
Sampson	1,764	0.7	7.9	11.6	62.0	17.0	0.7	0.0
Scotland	1,265	0.6	11.6	14.8	59.3	13.0	0.8	0.0
Stanly	1,160	0.6	9.6	16.6	59.4	12.7	1.1	0.0
Stokes	741	0.5	8.4	15.1	63.0	12.3	0.7	0.0
Surry	1,548	0.1	7.8	14.7	58.6	17.7	1.2	0.0
Swain	450	0.7	8.9	14.4	63.1	12.2	0.7	0.0
Transylvan	518	0.0	9.3	17.0	60.0	12.5	1.2	0.0
Tyrrell	127	0.0	9.4	15.7	57.5	16.5	0.8	0.0
Union	2,583	0.9	8.6	13.5	57.4	18.5	1.2	0.0
Vance	1,504	1.0	10.2	15.2	56.8	15.6	1.1	0.0
Wake	12,419	0.5	6.1	11.2	59.4	21.6	1.1	0.0
Warren	511	0.8	12.3	13.5	55.6	16.0	1.8	0.0
Washington	398	0.8	8.8	16.1	58.0	14.8	1.5	0.0
Watauga	530	0.6	6.4	9.6	64.2	18.7	0.6	0.0

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

Table 4B
2005 - 2007 Pregnancy Nutrition Surveillance ⁽¹⁾
North Carolina
Comparison of Age Distribution by County
Combined 3 Years

Comparison of Age Distribution ⁽²⁾								
County	Number Included in Analysis	< 15 Years %	15 - 17 Years %	18 - 19 Years %	20 - 29 Years %	30 - 39 Years %	>= 40 Years %	Unknown %
Wayne Coun	2,992	0.3	7.9	13.7	63.0	14.3	0.8	0.0
Wilkes	1,412	0.4	8.2	14.9	61.0	14.9	0.6	0.0
Wilson Cou	2,147	0.8	8.8	15.5	58.9	15.0	0.9	0.0
Yadkin	783	0.1	9.6	12.3	60.4	17.0	0.6	0.0
Yancey	315	0.0	4.4	12.7	64.8	17.8	0.3	0.0
Unknown Counties	946	1.6	10.8	17.1	56.2	13.8	0.4	0.0
North Carolina	186,585	0.5	7.8	14.0	60.5	16.3	0.9	0.0

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

Table 5B
2005 - 2007 Pregnancy Nutrition Surveillance ⁽¹⁾
North Carolina
Comparison of Maternal Health Indicators by County
Combined 3 Years

Comparison of Maternal Health Indicators ^(2, 3)																		
County	Prepregnancy BMI (4)			Weight Gain (5)			Anemia (Low Hb/Hct) (6)				Gestational Diabetes During Pregnancy		Hypertension During Pregnancy					
	Under-weight	Over-weight		< Ideal	> Ideal		3rd Trimester		Postpartum									
							Number	% (Rank)	%	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)		
Alamance	1,690	13.1 (68)	43.5 (22)	759	21.9 (15)	40.2 (45)	370	36.5 (33)	1,806	37.5 (61)	0	*	*	0	*	*		
Alexander	595	13.1 (68)	45.4 (33)	430	30.7 (74)	37.4 (24)	113	21.2 (5)	284	48.6 (76)	0	*	*	0	*	*		
Alleghany	125	8.0 (6)	56.8 (89)	104	26.9 (53)	39.4 (37)	15	*	*	57	*	*	0	*	*	0	*	*
Anson	584	13.7 (78)	49.7 (64)	497	27.6 (59)	42.1 (59)	82	*	*	133	35.3 (58)	0	*	*	0	*	*	
Ashe	405	13.6 (73)	50.1 (67)	364	26.1 (46)	38.5 (32)	70	*	*	136	19.1 (16)	0	*	*	0	*	*	
Avery	268	13.4 (72)	43.3 (21)	24	*	*	29	*	*	43	*	*	0	*	*	0	*	*
Beaufort	837	9.4 (17)	57.9 (92)	454	24.7 (34)	41.0 (52)	110	39.1 (40)	832	31.4 (47)	0	*	*	0	*	*		
Bertie	460	6.7 (2)	64.6 (98)	260	44.2 (91)	34.2 (12)	72	*	*	264	62.5 (84)	0	*	*	0	*	*	
Bladen	778	9.9 (28)	49.2 (60)	483	32.3 (83)	36.6 (20)	68	*	*	205	51.7 (81)	0	*	*	0	*	*	
Brunswick	858	9.1 (15)	64.2 (97)	321	17.1 (2)	50.2 (90)	61	*	*	621	13.7 (6)	0	*	*	0	*	*	
Buncombe	3,457	13.9 (80)	40.9 (12)	2,647	20.6 (8)	42.3 (61)	474	27.0 (15)	1,521	30.6 (45)	0	*	*	0	*	*		
Burke	1,276	12.7 (64)	43.0 (17)	1,095	26.8 (52)	36.5 (19)	277	27.8 (17)	648	27.0 (37)	0	*	*	0	*	*		
Cabarrus	2,795	10.8 (40)	44.9 (28)	2,312	28.5 (65)	40.5 (47)	346	37.3 (35)	825	21.3 (24)	0	*	*	0	*	*		
Caldwell	1,383	14.7 (87)	46.9 (45)	1,224	22.5 (18)	45.2 (77)	148	19.6 (3)	500	19.4 (18)	0	*	*	0	*	*		
Camden	79	*	*	54	*	*	10	*	*	75	*	*	0	*	*	0	*	*
Carteret	759	10.9 (42)	44.3 (26)	661	16.8 (1)	45.4 (80)	202	30.2 (20)	446	23.1 (27)	0	*	*	0	*	*		
Caswell	308	12.7 (64)	55.8 (85)	262	25.6 (41)	38.5 (32)	43	*	*	95	*	*	0	*	*	0	*	*

- (1) Reporting period is January 1 through December 31.
- (2) Excludes records with unknown data or errors.
- (3) Rank compares one county's rate to other counties. Rank 1 = best rate.
- (4) Based on 1990 IOM report, "Nutrition During Pregnancy": underweight (BMI <19.8), overweight (BMI >26.0; includes overweight and obese women).
- (5) Based on 1990 IOM report, "Nutrition During Pregnancy": underweight prepregnancy (ideal weight gain = 28 to 40 lbs), normal weight prepregnancy (ideal weight gain = 25 to 35 lbs), overweight prepregnancy (ideal weight gain = 15 to 25 lbs), obese prepregnancy (ideal weight gain = 15 to 25 lbs).
- (6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.
- * Percentages are not calculated if <100 records are available for analysis after exclusions.

Table 5B
2005 - 2007 Pregnancy Nutrition Surveillance ⁽¹⁾
North Carolina
Comparison of Maternal Health Indicators by County
Combined 3 Years

Comparison of Maternal Health Indicators ^(2, 3)																		
County	Prepregnancy BMI ⁽⁴⁾			Weight Gain ⁽⁵⁾			Anemia (Low Hb/Hct) ⁽⁶⁾				Gestational Diabetes During Pregnancy	Hypertension During Pregnancy						
	Under-weight	Over-weight		< Ideal	> Ideal		3rd Trimester		Postpartum									
							Number	% (Rank)	Number	% (Rank)			Number	% (Rank)	Number	% (Rank)		
Catawba	2,650	12.0 (53)	44.9 (28)	2,166	36.1 (88)	31.2 (3)	582	24.2 (10)	1,809	24.5 (31)	0	*	*	0	*	*		
Chatham	803	8.1 (7)	52.8 (78)	667	26.4 (48)	37.5 (25)	49	*	*	675	24.0 (29)	0	*	*	0	*	*	
Cherokee	423	12.8 (66)	48.7 (58)	371	28.3 (63)	44.5 (76)	94	*	*	143	12.6 (5)	0	*	*	0	*	*	
Chowan	297	9.8 (25)	55.9 (86)	227	27.3 (55)	36.6 (20)	36	*	*	283	40.6 (68)	0	*	*	0	*	*	
Clay	125	13.6 (73)	44.0 (23)	105	20.0 (4)	50.5 (91)	20	*	*	56	*	*	0	*	*	0	*	*
Cleveland	1,708	13.6 (73)	44.9 (28)	111	25.2 (37)	47.7 (88)	216	38.4 (37)	1,580	26.1 (34)	0	*	*	0	*	*		
Columbus	1,079	10.6 (38)	56.4 (88)	604	28.3 (63)	41.4 (53)	158	36.7 (34)	201	44.8 (75)	0	*	*	0	*	*		
Craven	1,573	14.9 (89)	45.6 (35)	981	23.6 (23)	41.6 (57)	77	*	*	1,698	25.6 (33)	0	*	*	0	*	*	
Cumberland	7,393	11.7 (52)	44.3 (26)	5,095	20.6 (8)	47.2 (87)	1,836	44.7 (44)	2,716	39.4 (66)	0	*	*	0	*	*		
Currituck	162	13.6 (73)	41.4 (13)	71	*	*	32	*	*	175	30.3 (43)	0	*	*	0	*	*	
Dare	403	14.6 (86)	38.0 (5)	272	25.4 (39)	34.6 (13)	49	*	*	223	21.1 (23)	0	*	*	0	*	*	
Davidson	2,169	15.3 (91)	42.5 (15)	1,689	26.4 (48)	39.4 (37)	393	25.2 (12)	1,286	18.6 (14)	0	*	*	0	*	*		
Davie	512	13.1 (68)	46.9 (45)	430	24.0 (29)	43.7 (72)	98	*	*	315	20.3 (21)	0	*	*	0	*	*	
Duplin	1,138	25.0 (96)	43.1 (18)	775	31.6 (79)	32.9 (6)	196	32.7 (25)	490	42.7 (70)	0	*	*	0	*	*		
Durham	3,611	10.4 (36)	45.7 (36)	2,977	30.5 (72)	34.1 (11)	110	34.5 (29)	332	27.7 (39)	0	*	*	0	*	*		
Edgecombe	748	46.8 (98)	35.2 (2)	448	41.1 (90)	26.1 (1)	175	45.7 (48)	761	34.4 (54)	0	*	*	0	*	*		
Forsyth	4,700	11.6 (49)	45.3 (32)	3,359	26.3 (47)	39.6 (39)	861	39.5 (41)	2,920	38.3 (64)	0	*	*	0	*	*		

- (1) Reporting period is January 1 through December 31.
- (2) Excludes records with unknown data or errors.
- (3) Rank compares one county's rate to other counties. Rank 1 = best rate.
- (4) Based on 1990 IOM report, "Nutrition During Pregnancy": underweight (BMI <19.8), overweight (BMI >26.0; includes overweight and obese women).
- (5) Based on 1990 IOM report, "Nutrition During Pregnancy": underweight prepregnancy (ideal weight gain = 28 to 40 lbs), normal weight prepregnancy (ideal weight gain = 25 to 35 lbs), overweight prepregnancy (ideal weight gain = 15 to 25 lbs), obese prepregnancy (ideal weight gain = 15 to 25 lbs).
- (6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.
- * Percentages are not calculated if <100 records are available for analysis after exclusions.

Table 5B
2005 - 2007 Pregnancy Nutrition Surveillance ⁽¹⁾
North Carolina
Comparison of Maternal Health Indicators by County
Combined 3 Years

Comparison of Maternal Health Indicators ^(2, 3)																
County	Prepregnancy BMI ⁽⁴⁾			Weight Gain ⁽⁵⁾			Anemia (Low Hb/Hct) ⁽⁶⁾				Gestational Diabetes During Pregnancy		Hypertension During Pregnancy			
	Under-weight	Over-weight		< Ideal	> Ideal		3rd Trimester		Postpartum							
	Number	% (Rank)	%	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)		
Franklin	858	7.9 (4)	50.2 (68)	708	30.4 (71)	35.3 (16)	138	33.3 (26)	210	18.6 (14)	0	*	*	0	*	*
Gaston	3,430	14.1 (82)	45.7 (36)	2,843	30.1 (68)	40.1 (44)	769	23.1 (9)	2,609	43.7 (74)	0	*	*	0	*	*
Gates	82	*	*	58	*	*	15	*	55	*	0	*	*	0	*	*
Graham	199	9.5 (19)	38.2 (6)	147	18.4 (3)	46.9 (86)	11	*	59	*	0	*	*	0	*	*
Granville	796	8.3 (9)	54.4 (84)	693	31.9 (81)	35.2 (15)	79	*	211	36.5 (59)	0	*	*	0	*	*
Greene	483	8.1 (7)	50.9 (70)	408	35.3 (87)	33.1 (7)	71	*	160	43.1 (72)	0	*	*	0	*	*
Guilford	6,088	11.2 (47)	47.4 (52)	3,399	24.9 (36)	41.5 (54)	651	33.3 (26)	2,250	21.6 (25)	0	*	*	0	*	*
Halifax	1,306	9.3 (16)	56.2 (87)	1,180	27.5 (57)	40.4 (46)	128	48.4 (51)	641	50.4 (80)	0	*	*	0	*	*
Harnett	1,935	9.6 (21)	48.7 (58)	1,403	25.4 (39)	41.8 (58)	423	38.3 (36)	849	33.5 (52)	0	*	*	0	*	*
Haywood	749	13.2 (71)	41.9 (14)	591	21.5 (13)	43.0 (66)	83	*	484	15.3 (8)	0	*	*	0	*	*
Henderson	971	9.8 (25)	46.1 (39)	513	22.4 (17)	42.1 (59)	231	27.3 (16)	634	29.2 (41)	0	*	*	0	*	*
Hertford	560	7.9 (4)	60.9 (96)	428	23.8 (26)	43.7 (72)	54	*	123	79.7 (86)	0	*	*	0	*	*
Hoke	779	8.9 (14)	52.8 (78)	570	21.9 (15)	43.2 (70)	85	*	392	33.4 (51)	0	*	*	0	*	*
Hyde	87	*	*	72	*	*	16	*	40	*	0	*	*	0	*	*
Iredell	2,371	12.6 (63)	46.9 (45)	2,135	23.9 (28)	43.1 (67)	564	26.4 (14)	2,390	12.4 (4)	0	*	*	0	*	*
Jackson	626	9.6 (21)	50.0 (65)	517	20.9 (12)	46.0 (82)	68	*	398	17.6 (12)	0	*	*	0	*	*
Johnston	1,562	10.2 (32)	50.0 (65)	874	26.5 (50)	38.6 (34)	487	35.3 (30)	1,722	31.7 (49)	0	*	*	0	*	*

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one county's rate to other counties. Rank 1 = best rate.
(4) Based on 1990 IOM report, "Nutrition During Pregnancy": underweight (BMI <19.8), overweight (BMI >26.0; includes overweight and obese women).
(5) Based on 1990 IOM report, "Nutrition During Pregnancy": underweight prepregnancy (ideal weight gain = 28 to 40 lbs), normal weight prepregnancy (ideal weight gain = 25 to 35 lbs), overweight prepregnancy (ideal weight gain = 15 to 25 lbs), obese prepregnancy (ideal weight gain = 15 to 25 lbs).
(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

Table 5B
2005 - 2007 Pregnancy Nutrition Surveillance ⁽¹⁾
North Carolina
Comparison of Maternal Health Indicators by County
Combined 3 Years

Comparison of Maternal Health Indicators ^(2, 3)														
County	Prepregnancy BMI ⁽⁴⁾			Weight Gain ⁽⁵⁾			Anemia (Low Hb/Hct) ⁽⁶⁾				Gestational Diabetes During Pregnancy	Hypertension During Pregnancy		
	Under-weight		Over-weight	< Ideal		> Ideal	3rd Trimester		Postpartum					
	Number	% (Rank)	%	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)			Number	% (Rank)
Jones	211	9.5 (19)	58.3 (94)	14	* *	* *	23	* *	22	* *	0	* *	0	* *
Lee	1,190	8.3 (9)	47.4 (52)	948	27.3 (55)	40.7 (48)	185	45.4 (47)	574	34.5 (55)	0	* *	0	* *
Lenoir	919	12.1 (54)	49.6 (63)	537	24.0 (29)	44.1 (75)	244	40.2 (42)	854	34.9 (56)	0	* *	0	* *
Lincoln	1,048	14.3 (84)	42.9 (16)	799	23.7 (25)	42.8 (64)	151	20.5 (4)	614	11.7 (3)	0	* *	0	* *
Macon	759	12.4 (60)	46.9 (45)	619	34.7 (86)	29.9 (2)	92	* *	386	24.1 (30)	0	* *	0	* *
Madison	594	9.8 (25)	43.1 (18)	359	23.4 (21)	40.9 (50)	87	* *	368	10.9 (1)	0	* *	0	* *
Martin	331	15.1 (90)	47.4 (52)	127	26.0 (44)	40.9 (50)	40	* *	77	* *	0	* *	0	* *
Mcdowell	511	11.5 (48)	57.5 (90)	419	30.5 (72)	37.5 (25)	77	* *	144	48.6 (76)	0	* *	0	* *
Mecklenbur	7,732	10.2 (32)	47.7 (55)	5,212	22.6 (20)	43.1 (67)	1,627	35.5 (31)	3,933	32.3 (50)	0	* *	0	* *
Mitchell	213	10.3 (35)	46.5 (40)	158	25.3 (38)	38.6 (34)	20	* *	111	14.4 (7)	0	* *	0	* *
Montgomery	540	10.0 (31)	49.4 (62)	276	27.5 (57)	37.7 (29)	84	* *	113	27.4 (38)	0	* *	0	* *
Moore	877	10.9 (42)	51.1 (73)	502	23.5 (22)	43.6 (71)	137	34.3 (28)	300	56.7 (83)	0	* *	0	* *
Nash	1,655	12.4 (60)	48.1 (56)	1,215	30.3 (70)	35.6 (17)	139	43.2 (43)	459	37.7 (62)	0	* *	0	* *
New Hanove	1,933	11.0 (45)	45.4 (33)	1,506	23.8 (26)	42.5 (62)	236	32.2 (24)	993	26.7 (36)	0	* *	0	* *
Northhampt	454	8.6 (12)	57.9 (92)	331	26.0 (44)	42.6 (63)	66	* *	164	42.7 (70)	0	* *	0	* *
Onslow	4,935	18.2 (94)	37.5 (3)	3,825	20.3 (6)	45.2 (77)	980	21.8 (8)	3,878	19.3 (17)	0	* *	0	* *
Orange	943	10.4 (36)	50.3 (69)	723	24.3 (32)	46.6 (85)	62	* *	618	23.3 (28)	0	* *	0	* *

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one county's rate to other counties. Rank 1 = best rate.
(4) Based on 1990 IOM report, "Nutrition During Pregnancy": underweight (BMI <19.8), overweight (BMI >26.0; includes overweight and obese women).
(5) Based on 1990 IOM report, "Nutrition During Pregnancy": underweight prepregnancy (ideal weight gain = 28 to 40 lbs), normal weight prepregnancy (ideal weight gain = 25 to 35 lbs), overweight prepregnancy (ideal weight gain = 15 to 25 lbs), obese prepregnancy (ideal weight gain = 15 to 25 lbs).
(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

Table 5B
2005 - 2007 Pregnancy Nutrition Surveillance ⁽¹⁾
North Carolina
Comparison of Maternal Health Indicators by County
Combined 3 Years

Comparison of Maternal Health Indicators ^(2, 3)														
County	Prepregnancy BMI (4)			Weight Gain (5)			Anemia (Low Hb/Hct) (6)				Gestational Diabetes During Pregnancy		Hypertension During Pregnancy	
	Under- weight		Over- weight	< Ideal		> Ideal	3rd Trimester		Postpartum					
	Number	%	%	Number	%	%	Number	%	Number	%	Number	%	Number	%
		(Rank)			(Rank)	(Rank)		(Rank)		(Rank)		(Rank)		(Rank)
Pamlico	204	12.3 (59)	47.1 (50)	62	* *	* *	17	* *	41	* *	0	* *	0	* *
Pasquotank	653	8.7 (13)	54.1 (82)	498	31.3 (76)	33.1 (7)	127	10.2 (1)	728	73.1 (85)	0	* *	0	* *
Pender	725	10.9 (42)	46.5 (40)	636	24.8 (35)	39.9 (42)	138	31.2 (22)	199	26.1 (34)	0	* *	0	* *
Perquimans	142	6.3 (1)	59.2 (95)	17	* *	* *	24	* *	131	48.9 (78)	0	* *	0	* *
Person	671	11.6 (49)	46.6 (42)	425	24.0 (29)	43.1 (67)	104	29.8 (19)	262	33.6 (53)	0	* *	0	* *
Pitt	2,837	9.9 (28)	53.8 (81)	2,312	26.6 (51)	38.8 (36)	548	48.2 (50)	717	42.0 (69)	0	* *	0	* *
Polk	141	19.1 (95)	37.6 (4)	68	* *	* *	15	* *	69	* *	0	* *	0	* *
Randolph	2,023	12.2 (57)	48.6 (57)	1,648	20.8 (11)	45.3 (79)	382	17.3 (2)	1,855	17.3 (10)	0	* *	0	* *
Richmond	1,151	17.6 (93)	40.0 (9)	929	26.9 (53)	44.0 (74)	235	48.9 (52)	572	37.1 (60)	0	* *	0	* *
Robeson	3,406	13.6 (73)	47.3 (51)	2,099	33.5 (85)	33.8 (10)	770	21.3 (6)	1,139	38.2 (63)	0	* *	0	* *
Rockingham	870	12.5 (62)	47.0 (49)	297	25.6 (41)	46.1 (83)	262	25.2 (12)	469	17.5 (11)	0	* *	0	* *
Rowan	2,350	14.0 (81)	44.1 (24)	2,019	28.0 (61)	39.7 (40)	534	36.1 (32)	1,225	31.0 (46)	0	* *	0	* *
Rutherford	727	37.7 (97)	32.2 (1)	189	38.6 (89)	32.3 (4)	49	* *	296	28.7 (40)	0	* *	0	* *
Sampson	1,209	7.5 (3)	54.3 (83)	907	24.5 (33)	46.5 (84)	247	45.3 (46)	683	30.5 (44)	0	* *	0	* *
Scotland	935	9.7 (23)	52.4 (76)	759	29.1 (66)	39.9 (42)	77	* *	176	43.2 (73)	0	* *	0	* *
Stanly	965	14.8 (88)	39.9 (8)	712	31.6 (79)	37.6 (28)	154	64.3 (53)	657	49.3 (79)	0	* *	0	* *
Stokes	500	16.2 (92)	45.2 (31)	398	20.1 (5)	49.7 (89)	111	30.6 (21)	171	39.8 (67)	0	* *	0	* *

- (1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one county's rate to other counties. Rank 1 = best rate.
(4) Based on 1990 IOM report, "Nutrition During Pregnancy": underweight (BMI <19.8), overweight (BMI >26.0; includes overweight and obese women).
(5) Based on 1990 IOM report, "Nutrition During Pregnancy": underweight prepregnancy (ideal weight gain = 28 to 40 lbs), normal weight prepregnancy (ideal weight gain = 25 to 35 lbs), overweight prepregnancy (ideal weight gain = 15 to 25 lbs), obese prepregnancy (ideal weight gain = 15 to 25 lbs).
(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

Table 5B
2005 - 2007 Pregnancy Nutrition Surveillance ⁽¹⁾
North Carolina
Comparison of Maternal Health Indicators by County
Combined 3 Years

Comparison of Maternal Health Indicators ^(2, 3)																		
County	Prepregnancy BMI ⁽⁴⁾			Weight Gain ⁽⁵⁾			Anemia (Low Hb/Hct) ⁽⁶⁾				Gestational Diabetes During Pregnancy		Hypertension During Pregnancy					
	Under-weight	Over-weight		< Ideal	> Ideal		3rd Trimester		Postpartum									
	Number	% (Rank)	%	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)				
Surry	938	13.0 (67)	43.1 (18)	757	21.7 (14)	45.4 (80)	209	21.5 (7)	499	34.9 (56)	0	*	*	0	*	*		
Swain	337	11.6 (49)	52.5 (77)	174	25.9 (43)	39.7 (40)	28	*	*	153	17.6 (12)	0	*	*	0	*	*	
Transylvan	412	13.8 (79)	40.8 (11)	142	20.4 (7)	41.5 (54)	63	*	*	96	*	*	0	*	*	0	*	*
Tyrrell	108	10.2 (32)	52.8 (78)	82	*	*	9	*	*	31	*	*	0	*	*	0	*	*
Union	1,798	12.2 (57)	44.1 (24)	877	27.7 (60)	36.3 (18)	400	38.5 (38)	952	21.8 (26)	0	*	*	0	*	*		
Vance	702	11.1 (46)	51.0 (71)	212	30.2 (69)	36.8 (22)	116	38.8 (39)	326	54.9 (82)	0	*	*	0	*	*		
Wake	7,637	9.7 (23)	46.6 (42)	5,649	31.5 (77)	33.6 (9)	1,250	28.7 (18)	3,758	19.6 (19)	0	*	*	0	*	*		
Warren	445	9.9 (28)	51.0 (71)	367	31.1 (75)	37.3 (23)	52	*	*	127	38.6 (65)	0	*	*	0	*	*	
Washington	351	8.5 (11)	57.5 (90)	296	32.8 (84)	37.8 (30)	40	*	*	72	*	*	0	*	*	0	*	*
Watauga	430	12.1 (54)	40.0 (9)	199	20.6 (8)	40.7 (48)	55	*	*	190	11.1 (2)	0	*	*	0	*	*	
Wayne Coun	2,140	14.4 (85)	49.3 (61)	1,005	30.0 (67)	38.3 (31)	203	46.8 (49)	1,210	30.2 (42)	0	*	*	0	*	*		
Wilkes	1,242	12.1 (54)	51.5 (74)	1,037	31.5 (77)	37.5 (25)	227	24.7 (11)	533	19.9 (20)	0	*	*	0	*	*		
Wilson Cou	1,830	10.6 (38)	51.9 (75)	1,270	32.0 (82)	35.0 (14)	355	45.1 (45)	825	24.8 (32)	0	*	*	0	*	*		
Yadkin	657	9.4 (17)	46.6 (42)	515	22.5 (18)	42.9 (65)	111	31.5 (23)	293	16.4 (9)	0	*	*	0	*	*		
Yancey	251	10.8 (40)	38.2 (6)	117	28.2 (62)	32.5 (5)	16	*	*	100	21.0 (22)	0	*	*	0	*	*	
Unknown Counties	715	14.1 (82)	45.7 (36)	491	23.6 (23)	41.5 (54)	84	*	*	369	31.4 (47)	0	*	*	0	*	*	
North Carolina	133,380	12.3	46.8	93,573	26.5	40.2	22,607	33.5	69,913	29.5	0	*	*	0	*	*		

- (1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one county's rate to other counties. Rank 1 = best rate.
(4) Based on 1990 IOM report, "Nutrition During Pregnancy": underweight (BMI <19.8), overweight (BMI >26.0; includes overweight and obese women).
(5) Based on 1990 IOM report, "Nutrition During Pregnancy": underweight prepregnancy (ideal weight gain = 28 to 40 lbs), normal weight prepregnancy (ideal weight gain = 25 to 35 lbs), overweight prepregnancy (ideal weight gain = 15 to 25 lbs), obese prepregnancy (ideal weight gain = 15 to 25 lbs).
(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

Table 6B
2005 - 2007 Pregnancy Nutrition Surveillance ⁽¹⁾
North Carolina
Comparison of Maternal Behavioral Indicators by County
Combined 3 Years

Comparison of Maternal Behavioral Indicators ^(2, 3)										
County	Medical Care			WIC Enrollment		Multivitamin Consumption				
	1st Trimester		No Care	1st Trimester		Prepregnancy (4)		During Pregnancy (5)		
	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	
Alamance	2,565	73.7 (72)	0.6 (32)	3,187	36.6 (42)	0	* *	0	* *	
Alexander	558	75.1 (67)	1.6 (89)	701	29.2 (68)	0	* *	0	* *	
Alleghany	134	87.3 (16)	0.0 (1)	152	42.1 (23)	0	* *	0	* *	
Anson	620	79.4 (45)	1.0 (67)	685	41.5 (26)	0	* *	0	* *	
Ashe	464	87.3 (16)	0.4 (19)	483	45.5 (14)	0	* *	0	* *	
Avery	65	* *	* *	308	44.8 (16)	0	* *	0	* *	
Beaufort	944	74.3 (71)	0.7 (45)	1,088	46.2 (13)	0	* *	0	* *	
Bertie	348	83.6 (33)	1.1 (76)	522	41.4 (27)	0	* *	0	* *	
Bladen	624	70.2 (80)	1.0 (67)	904	39.8 (34)	0	* *	0	* *	
Brunswick	1,379	79.0 (48)	0.6 (32)	1,772	34.7 (44)	0	* *	0	* *	
Buncombe	3,670	91.4 (2)	0.3 (12)	4,155	34.4 (45)	0	* *	0	* *	
Burke	1,666	65.9 (89)	0.5 (28)	1,704	18.5 (94)	0	* *	0	* *	
Cabarrus	3,300	68.6 (85)	0.6 (32)	3,372	27.2 (76)	0	* *	0	* *	
Caldwell	1,535	91.9 (1)	0.3 (12)	1,615	43.8 (20)	0	* *	0	* *	
Camden	93	* *	* *	101	32.7 (52)	0	* *	0	* *	
Carteret	961	85.2 (25)	0.6 (32)	1,023	22.8 (88)	0	* *	0	* *	
Caswell	340	79.1 (46)	0.6 (32)	369	45.5 (14)	0	* *	0	* *	
Catawba	2,588	74.4 (70)	0.7 (45)	2,954	25.6 (82)	0	* *	0	* *	
Chatham	899	75.5 (64)	0.8 (56)	975	30.1 (65)	0	* *	0	* *	
Cherokee	497	83.9 (32)	0.4 (19)	528	33.5 (48)	0	* *	0	* *	

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one county's rate to other counties. Rank 1 = best rate.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

Table 6B
2005 - 2007 Pregnancy Nutrition Surveillance ⁽¹⁾
North Carolina
Comparison of Maternal Behavioral Indicators by County
Combined 3 Years

Comparison of Maternal Behavioral Indicators ^(2, 3)									
County	Medical Care			WIC Enrollment		Multivitamin Consumption			
	1st Trimester		No Care	1st Trimester		Pregnancy (4)		During Pregnancy (5)	
	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Chowan	318	85.2 (25)	0.3 (12)	341	42.2 (22)	0	* *	0	* *
Clay	121	84.3 (29)	0.0 (1)	133	54.1 (2)	0	* *	0	* *
Cleveland	1,667	72.2 (75)	0.7 (45)	2,118	41.6 (25)	0	* *	0	* *
Columbus	853	63.1 (92)	1.4 (82)	1,310	31.8 (58)	0	* *	0	* *
Craven	2,147	80.3 (41)	0.0 (1)	2,626	32.8 (51)	0	* *	0	* *
Cumberland	8,987	83.5 (35)	0.8 (56)	9,673	23.0 (86)	0	* *	0	* *
Currituck	200	76.0 (59)	1.5 (84)	220	36.8 (39)	0	* *	0	* *
Dare	495	83.6 (33)	0.4 (19)	543	27.1 (77)	0	* *	0	* *
Davidson	2,667	74.7 (69)	0.4 (19)	2,896	27.1 (77)	0	* *	0	* *
Davie	572	87.8 (13)	0.5 (28)	611	30.6 (61)	0	* *	0	* *
Duplin	1,651	77.6 (52)	1.0 (67)	1,687	27.9 (75)	0	* *	0	* *
Durham	4,940	85.3 (24)	0.8 (56)	5,314	12.8 (99)	0	* *	0	* *
Edgecombe	1,578	64.5 (91)	2.3 (95)	1,632	32.2 (55)	0	* *	0	* *
Forsyth	6,476	78.4 (49)	0.7 (45)	7,000	17.6 (96)	0	* *	0	* *
Franklin	928	81.5 (39)	0.3 (12)	995	26.2 (79)	0	* *	0	* *
Gaston	3,997	55.0 (95)	0.9 (62)	4,222	16.7 (98)	0	* *	0	* *
Gates	77	* *	* *	90	* *	0	* *	0	* *
Graham	180	88.9 (10)	0.0 (1)	225	59.6 (1)	0	* *	0	* *
Granville	934	77.3 (54)	0.9 (62)	977	29.2 (68)	0	* *	0	* *
Greene	551	84.2 (30)	0.7 (45)	566	38.5 (36)	0	* *	0	* *

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one county's rate to other counties. Rank 1 = best rate.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

Table 6B
2005 - 2007 Pregnancy Nutrition Surveillance ⁽¹⁾
North Carolina
Comparison of Maternal Behavioral Indicators by County
Combined 3 Years

Comparison of Maternal Behavioral Indicators ^(2, 3)										
County	Medical Care			WIC Enrollment		Multivitamin Consumption				
	1st Trimester		No Care	1st Trimester		Prepregnancy (4)		During Pregnancy (5)		
	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	% (Rank)
Guilford	7,523	75.7 (61)	1.1 (76)	8,578	23.3 (84)	0	* *	0	* *	*
Halifax	1,501	86.1 (21)	0.6 (32)	1,587	26.0 (80)	0	* *	0	* *	*
Harnett	1,961	69.4 (83)	1.0 (67)	2,346	30.1 (65)	0	* *	0	* *	*
Haywood	807	86.4 (20)	0.2 (8)	920	40.8 (31)	0	* *	0	* *	*
Henderson	1,136	90.7 (5)	0.3 (12)	1,464	20.5 (92)	0	* *	0	* *	*
Hertford	534	86.0 (22)	0.2 (8)	648	29.5 (67)	0	* *	0	* *	*
Hoke	953	75.4 (66)	1.4 (82)	1,292	24.1 (83)	0	* *	0	* *	*
Hyde	96	* *	* *	106	36.8 (39)	0	* *	0	* *	*
Iredell	2,694	61.2 (93)	1.6 (89)	2,850	23.0 (86)	0	* *	0	* *	*
Jackson	719	90.0 (7)	0.7 (45)	775	48.9 (9)	0	* *	0	* *	*
Johnston	2,617	79.6 (44)	0.8 (56)	2,993	17.2 (97)	0	* *	0	* *	*
Jones	37	* *	* *	227	51.5 (4)	0	* *	0	* *	*
Lee	1,363	65.1 (90)	0.2 (8)	1,494	23.2 (85)	0	* *	0	* *	*
Lenoir	1,021	79.8 (43)	0.6 (32)	1,428	33.1 (50)	0	* *	0	* *	*
Lincoln	1,081	70.1 (81)	0.5 (28)	1,265	36.8 (39)	0	* *	0	* *	*
Macon	904	84.1 (31)	0.1 (7)	987	42.1 (23)	0	* *	0	* *	*
Madison	473	85.4 (23)	0.6 (32)	679	40.9 (29)	0	* *	0	* *	*
Martin	176	90.9 (4)	0.6 (32)	376	43.9 (18)	0	* *	0	* *	*
Mcdowell	548	76.1 (57)	1.5 (84)	605	41.2 (28)	0	* *	0	* *	*
Mecklenbur	10,111	68.5 (86)	1.6 (89)	12,606	9.2(100)	0	* *	0	* *	*

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one county's rate to other counties. Rank 1 = best rate.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

Table 6B
2005 - 2007 Pregnancy Nutrition Surveillance ⁽¹⁾
North Carolina
Comparison of Maternal Behavioral Indicators by County
Combined 3 Years

Comparison of Maternal Behavioral Indicators ^(2, 3)										
County	Medical Care				WIC Enrollment		Multivitamin Consumption			
	1st Trimester		No Care		1st Trimester		Prepregnancy (4)		During Pregnancy (5)	
	Number	% (Rank)	% (Rank)		Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Mitchell	224	90.2 (6)	0.4 (19)		251	46.6 (11)	0	* *	0	* *
Montgomery	449	71.3 (76)	1.6 (89)		725	39.6 (35)	0	* *	0	* *
Moore	927	76.9 (56)	1.0 (67)		1,209	28.8 (73)	0	* *	0	* *
Nash	1,818	70.3 (79)	1.5 (84)		2,043	32.6 (53)	0	* *	0	* *
New Hanove	2,924	70.6 (78)	1.5 (84)		3,253	29.0 (71)	0	* *	0	* *
Northhampt	446	87.0 (18)	0.7 (45)		532	30.5 (62)	0	* *	0	* *
Onslow	5,194	89.6 (9)	0.2 (8)		6,186	29.1 (70)	0	* *	0	* *
Orange	1,157	75.1 (67)	0.9 (62)		1,351	25.8 (81)	0	* *	0	* *
Pamlico	95	* *	* *		232	48.7 (10)	0	* *	0	* *
Pasquotank	811	73.4 (73)	1.0 (67)		893	30.7 (60)	0	* *	0	* *
Pender	886	71.2 (77)	1.2 (79)		920	30.5 (62)	0	* *	0	* *
Perquimans	191	79.1 (46)	0.0 (1)		199	31.7 (59)	0	* *	0	* *
Person	541	83.2 (37)	1.3 (80)		769	37.3 (38)	0	* *	0	* *
Pitt	3,121	84.5 (27)	0.9 (62)		3,418	31.9 (57)	0	* *	0	* *
Polk	116	82.8 (38)	1.7 (93)		196	33.7 (47)	0	* *	0	* *
Randolph	2,517	77.5 (53)	0.6 (32)		2,745	28.1 (74)	0	* *	0	* *
Richmond	1,352	75.9 (60)	1.0 (67)		1,469	35.3 (43)	0	* *	0	* *
Robeson	3,237	70.1 (81)	0.6 (32)		4,290	20.9 (90)	0	* *	0	* *
Rockingham	1,781	79.9 (42)	0.7 (45)		1,918	40.9 (29)	0	* *	0	* *
Rowan	2,712	68.0 (87)	1.5 (84)		2,836	21.5 (89)	0	* *	0	* *

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one county's rate to other counties. Rank 1 = best rate.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

Table 6B
2005 - 2007 Pregnancy Nutrition Surveillance ⁽¹⁾
North Carolina
Comparison of Maternal Behavioral Indicators by County
Combined 3 Years

Comparison of Maternal Behavioral Indicators ^(2, 3)										
County	Medical Care			WIC Enrollment		Multivitamin Consumption				
	1st Trimester		No Care	1st Trimester		Prepregnancy (4)		During Pregnancy (5)		
	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	% (Rank)
Rutherford	461	83.3 (36)	0.4 (19)	1,040	44.0 (17)	0	* *	0	* *	*
Sampson	1,616	75.6 (63)	0.6 (32)	1,726	32.0 (56)	0	* *	0	* *	*
Scotland	1,123	75.5 (64)	1.3 (80)	1,232	51.3 (5)	0	* *	0	* *	*
Stanly	940	77.1 (55)	0.3 (12)	1,143	29.0 (71)	0	* *	0	* *	*
Stokes	670	88.7 (11)	0.7 (45)	719	39.9 (33)	0	* *	0	* *	*
Surry	1,401	78.3 (50)	1.1 (76)	1,521	34.4 (45)	0	* *	0	* *	*
Swain	307	86.6 (19)	0.7 (45)	442	50.0 (7)	0	* *	0	* *	*
Transylvan	244	89.8 (8)	0.4 (19)	516	50.6 (6)	0	* *	0	* *	*
Tyrrell	111	91.0 (3)	0.0 (1)	121	53.7 (3)	0	* *	0	* *	*
Union	1,755	72.8 (74)	1.7 (93)	2,540	18.7 (93)	0	* *	0	* *	*
Vance	1,134	69.1 (84)	0.5 (28)	1,461	20.9 (90)	0	* *	0	* *	*
Wake	11,136	60.9 (94)	1.0 (67)	12,090	18.2 (95)	0	* *	0	* *	*
Warren	436	75.7 (61)	0.9 (62)	497	43.7 (21)	0	* *	0	* *	*
Washington	357	80.4 (40)	0.8 (56)	392	43.9 (18)	0	* *	0	* *	*
Watauga	321	88.2 (12)	0.3 (12)	524	49.8 (8)	0	* *	0	* *	*
Wayne Coun	1,696	66.0 (88)	1.0 (67)	2,947	30.2 (64)	0	* *	0	* *	*
Wilkes	1,234	84.4 (28)	0.4 (19)	1,403	33.2 (49)	0	* *	0	* *	*
Wilson Cou	1,897	78.1 (51)	0.8 (56)	2,105	32.3 (54)	0	* *	0	* *	*
Yadkin	726	87.6 (14)	0.4 (19)	775	40.5 (32)	0	* *	0	* *	*
Yancey	169	87.6 (14)	0.6 (32)	313	46.3 (12)	0	* *	0	* *	*

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one county's rate to other counties. Rank 1 = best rate.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

Table 6B
2005 - 2007 Pregnancy Nutrition Surveillance ⁽¹⁾
North Carolina
Comparison of Maternal Behavioral Indicators by County
Combined 3 Years

Comparison of Maternal Behavioral Indicators ^(2, 3)									
County	Medical Care			WIC Enrollment		Multivitamin Consumption			
	1st Trimester		No Care	1st Trimester		Pregpregnancy (4)		During Preganancy (5)	
	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Unknown Counties	750	76.1 (57)	0.7 (45)	925	38.4 (37)	0	* *	0	* *
North Carolina	157,827	75.6	0.9	182,870	26.8	0	*	0	*

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one county's rate to other counties. Rank 1 = best rate.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

Table 7B
2005 - 2007 Pregnancy Nutrition Surveillance (1)
North Carolina
Comparison of Smoking Indicators by County
Combined 3 Years

Comparison of Smoking Indicators (2, 3)										
County	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)		Smoking in Household			
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy				Prenatal		Postpartum	
	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Alamance	2,750	32.1 (58)	2,610	20.1 (54)	648	31.5 (26)	0	* *	0	* *
Alexander	595	41.0 (82)	556	30.0 (85)	165	17.0 (62)	0	* *	0	* *
Alleghany	123	30.9 (54)	134	24.6 (70)	29	* *	0	* *	0	* *
Anson	586	36.3 (73)	628	20.2 (55)	189	34.4 (17)	0	* *	0	* *
Ashe	402	50.5 (92)	463	33.3 (91)	186	24.2 (52)	0	* *	0	* *
Avery	269	28.6 (47)	67	* *	6	* *	0	* *	0	* *
Beaufort	905	32.7 (61)	932	17.4 (39)	239	36.4 (13)	0	* *	0	* *
Bertie	460	15.4 (8)	351	9.1 (5)	43	* *	0	* *	0	* *
Bladen	799	19.6 (24)	630	20.0 (52)	98	* *	0	* *	0	* *
Brunswick	1,271	36.0 (71)	1,382	27.0 (77)	351	29.3 (32)	0	* *	0	* *
Buncombe	3,568	32.6 (60)	3,671	20.8 (59)	995	27.0 (43)	0	* *	0	* *
Burke	725	56.4 (95)	1,668	26.4 (73)	381	19.7 (58)	0	* *	0	* *
Cabarrus	2,919	23.7 (34)	3,297	14.4 (24)	625	36.5 (12)	0	* *	0	* *
Caldwell	1,367	37.5 (76)	1,536	28.8 (82)	467	14.6 (63)	0	* *	0	* *
Camden	83	* *	93	* *	23	* *	0	* *	0	* *

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one county's rate to other counties. Rank 1 = best rate.

(4) Among women who smoked 3 months prior to pregnancy.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

Table 7B
2005 - 2007 Pregnancy Nutrition Surveillance (1)
North Carolina
Comparison of Smoking Indicators by County
Combined 3 Years

Comparison of Smoking Indicators (2, 3)										
County	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)		Smoking in Household			
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy				Prenatal		Postpartum	
	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Carteret	746	47.2 (89)	957	26.9 (76)	305	36.7 (11)	0	* *	0	* *
Caswell	317	35.6 (67)	342	27.5 (79)	102	21.6 (57)	0	* *	0	* *
Catawba	2,633	33.1 (62)	2,585	20.3 (56)	697	31.4 (27)	0	* *	0	* *
Chatham	834	12.6 (4)	902	13.4 (18)	80	* *	0	* *	0	* *
Cherokee	447	52.8 (93)	501	39.3 (95)	222	24.3 (51)	0	* *	0	* *
Chowan	309	35.9 (68)	319	16.9 (36)	99	* *	0	* *	0	* *
Clay	125	36.8 (75)	123	33.3 (91)	43	* *	0	* *	0	* *
Cleveland	1,763	42.5 (86)	1,669	22.9 (64)	541	29.2 (33)	0	* *	0	* *
Columbus	1,096	26.6 (42)	876	23.7 (67)	171	14.0 (64)	0	* *	0	* *
Craven	1,778	29.7 (49)	2,164	15.1 (27)	380	35.3 (15)	0	* *	0	* *
Cumberland	7,043	29.4 (48)	8,970	12.7 (14)	1,703	45.6 (3)	0	* *	0	* *
Currituck	193	74.1 (97)	197	29.4 (84)	126	4.0 (70)	0	* *	0	* *
Dare	391	32.2 (59)	487	18.9 (48)	100	37.0 (10)	0	* *	0	* *
Davidson	2,393	39.6 (79)	2,677	26.6 (74)	833	26.8 (44)	0	* *	0	* *
Davie	509	31.0 (55)	574	23.9 (68)	140	8.6 (68)	0	* *	0	* *

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one county's rate to other counties. Rank 1 = best rate.

(4) Among women who smoked 3 months prior to pregnancy.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

Table 7B
2005 - 2007 Pregnancy Nutrition Surveillance (1)
North Carolina
Comparison of Smoking Indicators by County
Combined 3 Years

Comparison of Smoking Indicators (2, 3)										
County	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)		Smoking in Household			
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy				Prenatal		Postpartum	
	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Duplin	1,363	13.1 (6)	1,652	9.0 (4)	166	28.3 (38)	0	* *	0	* *
Durham	3,715	14.7 (7)	4,959	6.1 (1)	459	47.3 (2)	0	* *	0	* *
Edgecombe	907	17.6 (16)	1,588	16.5 (33)	149	22.1 (56)	0	* *	0	* *
Forsyth	5,208	15.6 (9)	6,481	12.0 (9)	676	29.7 (29)	0	* *	0	* *
Franklin	851	26.3 (41)	933	12.3 (10)	195	35.4 (14)	0	* *	0	* *
Gaston	3,442	36.1 (72)	3,950	22.1 (62)	1,070	32.4 (24)	0	* *	0	* *
Gates	80	* *	74	* *	17	* *	0	* *	0	* *
Graham	197	33.5 (63)	181	32.6 (90)	46	* *	0	* *	0	* *
Granville	812	29.9 (51)	930	17.1 (37)	215	40.5 (7)	0	* *	0	* *
Greene	504	18.7 (20)	559	13.4 (18)	87	* *	0	* *	0	* *
Guilford	6,265	17.0 (12)	7,506	12.4 (11)	846	32.9 (23)	0	* *	0	* *
Halifax	1,371	27.3 (45)	1,518	15.9 (31)	351	43.6 (4)	0	* *	0	* *
Harnett	1,972	30.4 (53)	1,984	18.7 (47)	462	31.6 (25)	0	* *	0	* *
Haywood	741	53.8 (94)	807	32.0 (89)	330	29.1 (35)	0	* *	0	* *
Henderson	886	17.7 (18)	1,138	15.7 (28)	97	* *	0	* *	0	* *

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one county's rate to other counties. Rank 1 = best rate.

(4) Among women who smoked 3 months prior to pregnancy.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

Table 7B
2005 - 2007 Pregnancy Nutrition Surveillance (1)
North Carolina
Comparison of Smoking Indicators by County
Combined 3 Years

Comparison of Smoking Indicators (2, 3)										
County	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)		Smoking in Household			
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy				Prenatal		Postpartum	
	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Hertford	569	21.3 (28)	537	9.9 (6)	90	* *	0	* *	0	* *
Hoke	1,049	21.4 (29)	956	12.6 (12)	153	41.2 (5)	0	* *	0	* *
Hyde	88	* *	96	* *	18	* *	0	* *	0	* *
Iredell	2,380	29.8 (50)	2,701	21.6 (60)	647	25.3 (47)	0	* *	0	* *
Jackson	565	35.0 (65)	718	24.2 (69)	177	18.1 (60)	0	* *	0	* *
Johnston	1,937	18.7 (20)	2,624	13.3 (17)	274	33.9 (19)	0	* *	0	* *
Jones	211	41.2 (83)	39	* *	10	* *	0	* *	0	* *
Lee	1,230	27.2 (44)	1,398	15.7 (28)	285	34.4 (17)	0	* *	0	* *
Lenoir	1,231	19.3 (23)	1,058	17.3 (38)	138	29.0 (36)	0	* *	0	* *
Lincoln	1,094	41.6 (84)	1,078	27.8 (80)	376	30.1 (28)	0	* *	0	* *
Macon	777	35.9 (68)	931	27.2 (78)	248	19.0 (59)	0	* *	0	* *
Madison	603	42.5 (86)	475	23.4 (65)	150	33.3 (22)	0	* *	0	* *
Martin	301	43.2 (88)	177	30.5 (87)	45	* *	0	* *	0	* *
Mcdowell	523	22.4 (32)	555	15.9 (31)	101	29.7 (29)	0	* *	0	* *
Mecklenbur	8,606	5.4 (1)	10,179	7.7 (3)	343	39.1 (8)	0	* *	0	* *

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one county's rate to other counties. Rank 1 = best rate.

(4) Among women who smoked 3 months prior to pregnancy.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

Table 7B
2005 - 2007 Pregnancy Nutrition Surveillance (1)
North Carolina
Comparison of Smoking Indicators by County
Combined 3 Years

Comparison of Smoking Indicators (2, 3)										
County	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)		Smoking in Household			
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy				Prenatal		Postpartum	
	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Mitchell	225	50.2 (91)	228	35.1 (93)	100	25.0 (48)	0	* *	0	* *
Montgomery	616	25.6 (40)	457	14.9 (26)	86	* *	0	* *	0	* *
Moore	882	22.3 (31)	936	15.8 (30)	137	27.7 (41)	0	* *	0	* *
Nash	1,723	20.7 (26)	1,855	16.5 (33)	298	24.5 (50)	0	* *	0	* *
New Hanove	1,712	24.0 (35)	2,931	17.8 (44)	347	40.6 (6)	0	* *	0	* *
Northhampt	453	20.3 (25)	449	17.6 (42)	78	* *	0	* *	0	* *
Onslow	5,127	33.5 (63)	5,212	13.0 (15)	1,336	53.5 (1)	0	* *	0	* *
Orange	1,075	16.6 (11)	1,150	17.4 (39)	131	11.5 (65)	0	* *	0	* *
Pamlico	212	36.3 (73)	99	* *	32	* *	0	* *	0	* *
Pasquotank	689	17.0 (12)	812	11.5 (7)	92	* *	0	* *	0	* *
Pender	736	24.0 (35)	889	20.5 (58)	165	17.6 (61)	0	* *	0	* *
Perquimans	171	18.7 (20)	191	16.8 (35)	31	* *	0	* *	0	* *
Person	685	40.4 (80)	535	20.0 (52)	169	33.7 (20)	0	* *	0	* *
Pitt	2,837	21.1 (27)	3,114	12.6 (12)	498	39.0 (9)	0	* *	0	* *
Polk	100	28.0 (46)	117	19.7 (50)	15	* *	0	* *	0	* *

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one county's rate to other counties. Rank 1 = best rate.

(4) Among women who smoked 3 months prior to pregnancy.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

Table 7B
2005 - 2007 Pregnancy Nutrition Surveillance (1)
North Carolina
Comparison of Smoking Indicators by County
Combined 3 Years

Comparison of Smoking Indicators (2, 3)										
County	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)		Smoking in Household			
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy				Prenatal		Postpartum	
	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Randolph	2,168	31.8 (56)	2,511	18.4 (46)	566	27.2 (42)	0	* *	0	* *
Richmond	1,252	35.9 (68)	1,372	22.0 (61)	394	34.8 (16)	0	* *	0	* *
Robeson	3,422	17.6 (16)	3,272	19.9 (51)	398	7.8 (69)	0	* *	0	* *
Rockingham	1,651	38.4 (78)	1,799	25.1 (72)	569	27.9 (40)	0	* *	0	* *
Rowan	1,792	35.2 (66)	2,727	20.3 (56)	573	28.4 (37)	0	* *	0	* *
Rutherford	582	38.3 (77)	461	24.7 (71)	76	* *	0	* *	0	* *
Sampson	1,393	15.9 (10)	1,624	11.5 (7)	179	29.6 (31)	0	* *	0	* *
Scotland	1,130	24.1 (37)	1,129	23.5 (66)	237	11.0 (66)	0	* *	0	* *
Stanly	970	22.2 (30)	950	22.8 (63)	164	26.2 (45)	0	* *	0	* *
Stokes	535	56.6 (96)	669	37.1 (94)	267	23.2 (54)	0	* *	0	* *
Surry	1,177	32.0 (57)	1,402	28.3 (81)	320	10.9 (67)	0	* *	0	* *
Swain	321	77.3 (98)	307	26.7 (75)	137	24.1 (53)	0	* *	0	* *
Transylvan	432	49.8 (90)	243	29.2 (83)	79	* *	0	* *	0	* *
Tyrrell	109	18.3 (19)	107	14.0 (22)	15	* *	0	* *	0	* *
Union	1,441	13.0 (5)	1,761	14.7 (25)	89	* *	0	* *	0	* *

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one county's rate to other counties. Rank 1 = best rate.

(4) Among women who smoked 3 months prior to pregnancy.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

Table 7B
2005 - 2007 Pregnancy Nutrition Surveillance (1)
North Carolina
Comparison of Smoking Indicators by County
Combined 3 Years

Comparison of Smoking Indicators (2, 3)										
County	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)	Smoking in Household				
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy			Prenatal		Postpartum		
	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Vance	1,218	9.9 (2)	1,155	17.8 (44)	86	* *	0	* *	0	* *
Wake	7,453	11.1 (3)	11,326	7.3 (2)	700	26.1 (46)	0	* *	0	* *
Warren	445	17.5 (14)	444	13.1 (16)	63	* *	0	* *	0	* *
Washington	358	23.2 (33)	353	13.9 (21)	75	* *	0	* *	0	* *
Watauga	423	41.8 (85)	321	17.4 (39)	86	* *	0	* *	0	* *
Wayne Coun	2,641	24.2 (38)	1,706	14.2 (23)	300	33.7 (20)	0	* *	0	* *
Wilkes	1,241	40.5 (81)	1,235	30.2 (86)	427	23.0 (55)	0	* *	0	* *
Wilson Cou	1,924	17.5 (14)	1,900	13.4 (18)	292	28.1 (39)	0	* *	0	* *
Yadkin	662	30.2 (52)	729	19.2 (49)	178	29.2 (33)	0	* *	0	* *
Yancey	260	25.4 (39)	172	30.8 (88)	27	* *	0	* *	0	* *
Unknown Counties	852	26.6 (42)	755	17.6 (42)	166	24.7 (49)	0	* *	0	* *
North Carolina	139,972	25.4	158,548	16.3	28,186	30.8	0	*	0	*

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one county's rate to other counties. Rank 1 = best rate.

(4) Among women who smoked 3 months prior to pregnancy.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 8B
2005 - 2007 Pregnancy Nutrition Surveillance ⁽¹⁾
North Carolina
Comparison of Infant Health Indicators by County
Combined 3 Years

Comparison of Infant Health Indicators ^(2, 3)									
County	Birthweight			Preterm ⁽⁶⁾		Full Term LBW ⁽⁷⁾		Ever Breastfed	
	Low ⁽⁴⁾	High ⁽⁵⁾		Number	%	Number	%	Number	%
	Number	% ^(Rank)	% ^(Rank)						
Alamance	2,628	7.9 (20)	8.6 (88)	2,596	12.8 (27)	2,263	3.0 (11)	2,618	55.9 (51)
Alexander	555	9.2 (45)	7.9 (77)	546	11.5 (13)	483	5.6 (87)	312	87.2 (5)
Alleghany	134	8.2 (24)	3.0 (2)	134	14.9 (60)	114	3.5 (25)	119	78.2 (12)
Anson	625	11.2 (81)	5.4 (21)	612	17.2 (75)	507	4.5 (57)	622	35.2 (85)
Ashe	463	8.6 (31)	3.7 (4)	459	12.2 (17)	403	4.0 (42)	445	55.5 (52)
Avery	68	* * *	* *	65	* *	54	* *	57	* *
Beaufort	940	10.5 (73)	6.5 (52)	918	15.3 (63)	778	4.9 (69)	935	49.9 (66)
Bertie	347	14.4 (95)	6.1 (44)	341	18.5 (86)	278	6.1 (90)	346	20.8 (91)
Bladen	628	8.9 (39)	5.7 (26)	622	15.8 (66)	524	4.6 (60)	626	48.6 (68)
Brunswick	1,383	9.3 (47)	7.4 (69)	1,376	13.3 (31)	1,193	4.7 (64)	1,366	44.2 (75)
Buncombe	3,667	9.9 (60)	8.0 (81)	3,640	13.5 (34)	3,148	4.0 (42)	3,622	78.6 (11)
Burke	1,665	9.0 (41)	5.5 (23)	1,647	16.1 (71)	1,381	5.0 (74)	1,563	62.6 (30)
Cabarrus	3,332	9.2 (45)	5.9 (35)	3,272	13.8 (38)	2,821	3.7 (30)	3,328	66.3 (24)
Caldwell	1,536	8.7 (36)	6.9 (56)	1,521	12.4 (20)	1,332	5.4 (84)	1,529	51.7 (61)

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one county's rate to other counties. Rank 1 = best rate.
(4) <2500 g.
(5) >4000 g.
(6) <37 weeks gestation.
(7) 37 or more weeks gestation and <2500 g.
* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 8B
2005 - 2007 Pregnancy Nutrition Surveillance ⁽¹⁾
North Carolina
Comparison of Infant Health Indicators by County
Combined 3 Years

Comparison of Infant Health Indicators ^(2, 3)										
County	Birthweight				Preterm (6)		Full Term LBW (7)		Ever Breastfed	
	Low (4)		High (5)		Number	%	Number	%	Number	%
	Number	% (Rank)	% (Rank)	Number						
Camden	93	* * * *			93	* *			82	* *
Carteret	962	6.5 (7)	8.9 (89)		947	11.0 (9)			842	3.3 (22)
Caswell	342	12.9 (92)	5.3 (18)		340	14.4 (50)			291	6.5 (92)
Catawba	2,591	8.4 (29)	7.5 (71)		2,553	11.9 (15)			2,249	4.1 (47)
Chatham	915	6.0 (5)	8.4 (85)		893	11.8 (14)			788	1.9 (3)
Cherokee	502	9.6 (55)	6.2 (46)		499	17.2 (75)			413	3.6 (28)
Chowan	322	10.9 (78)	5.9 (35)		313	17.3 (80)			259	5.4 (84)
Clay	123	4.9 (3)	5.7 (26)		123	9.8 (5)			111	3.6 (28)
Cleveland	1,684	8.6 (31)	7.0 (60)		1,663	13.9 (40)			1,432	3.7 (30)
Columbus	893	11.9 (84)	6.0 (40)		868	19.8 (89)			695	3.7 (30)
Craven	2,174	7.8 (18)	8.3 (84)		2,123	11.3 (11)			1,883	3.8 (35)
Cumberland	8,984	9.9 (60)	7.4 (69)		8,886	16.1 (71)			7,455	4.5 (57)
Currituck	201	4.0 (1)	8.0 (81)		200	9.5 (4)			181	0.6 (1)
Dare	496	4.4 (2)	9.1 (90)		492	6.7 (1)			459	2.4 (5)

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one county's rate to other counties. Rank 1 = best rate.
(4) <2500 g.
(5) >4000 g.
(6) <37 weeks gestation.
(7) 37 or more weeks gestation and <2500 g.
* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 8B
2005 - 2007 Pregnancy Nutrition Surveillance ⁽¹⁾
North Carolina
Comparison of Infant Health Indicators by County
Combined 3 Years

Comparison of Infant Health Indicators ^(2, 3)										
County	Birthweight				Preterm (6)		Full Term LBW (7)		Ever Breastfed	
	Low (4)		High (5)							
	Number	% (Rank)	%	(Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Davidson	2,684	10.2 (66)	6.0	(40)	2,621	14.6 (55)	2,239	4.6 (60)	2,626	60.7 (34)
Davie	575	8.3 (27)	7.3	(67)	570	17.2 (75)	472	4.0 (42)	552	60.0 (39)
Duplin	1,653	8.0 (21)	6.8	(55)	1,631	12.5 (23)	1,427	3.2 (18)	1,615	56.0 (50)
Durham	5,003	10.3 (67)	6.6	(53)	4,931	17.9 (82)	4,048	5.3 (82)	4,843	65.8 (25)
Edgecombe	1,584	11.8 (83)	4.1	(5)	1,551	17.2 (75)	1,285	5.4 (84)	1,570	33.1 (88)
Forsyth	6,489	10.7 (76)	6.0	(40)	6,382	18.3 (85)	5,210	4.3 (54)	6,428	76.4 (16)
Franklin	940	10.3 (67)	7.6	(75)	933	16.0 (70)	784	6.3 (91)	928	47.6 (70)
Gaston	4,002	10.5 (73)	5.9	(35)	3,898	14.6 (55)	3,327	4.7 (64)	3,994	43.9 (76)
Gates	77	* * *	*	*	74	* *	61	* *	77	* *
Graham	181	8.3 (27)	9.4	(92)	180	10.0 (6)	162	4.3 (54)	48	* *
Granville	940	11.5 (82)	4.1	(5)	923	18.2 (84)	755	4.9 (69)	913	47.6 (70)
Greene	557	10.1 (65)	5.7	(26)	545	14.5 (53)	466	3.9 (38)	548	53.3 (58)
Guilford	7,532	9.9 (60)	5.8	(32)	7,409	14.1 (44)	6,368	3.9 (38)	7,474	65.7 (27)
Halifax	1,512	12.5 (89)	5.0	(12)	1,492	21.3 (93)	1,174	4.8 (67)	1,497	33.9 (87)

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one county's rate to other counties. Rank 1 = best rate.
(4) <2500 g.
(5) >4000 g.
(6) <37 weeks gestation.
(7) 37 or more weeks gestation and <2500 g.
* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 8B
2005 - 2007 Pregnancy Nutrition Surveillance ⁽¹⁾
North Carolina
Comparison of Infant Health Indicators by County
Combined 3 Years

Comparison of Infant Health Indicators ^(2, 3)										
County	Birthweight				Preterm (6)		Full Term LBW (7)		Ever Breastfed	
	Low (4)		High (5)							
	Number	% (Rank)	%	(Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Harnett	1,995	9.3 (47)	7.0 (60)		1,960	15.2 (62)	1,663	4.1 (47)	1,957	51.3 (63)
Haywood	806	10.4 (71)	7.2 (66)		801	14.9 (60)	682	4.1 (47)	695	60.6 (36)
Henderson	1,143	6.6 (8)	7.5 (71)		1,120	12.5 (23)	980	3.2 (18)	1,118	83.8 (8)
Hertford	534	10.3 (67)	5.2 (15)		529	20.2 (91)	422	3.8 (35)	530	38.1 (81)
Hoke	957	8.9 (39)	6.3 (49)		939	15.8 (66)	791	3.2 (18)	954	54.7 (54)
Hyde	96	* * * *			96	* *	80	* *	95	* *
Iredell	2,705	10.0 (63)	6.3 (49)		2,686	14.6 (55)	2,295	4.4 (56)	2,678	58.7 (44)
Jackson	717	6.7 (9)	9.9 (94)		709	11.4 (12)	628	2.9 (9)	703	69.3 (22)
Johnston	2,620	8.2 (24)	6.0 (40)		2,604	13.9 (40)	2,242	3.5 (25)	2,581	48.4 (69)
Jones	39	* * * *			35	* *	26	* *	39	* *
Lee	1,403	9.0 (41)	5.0 (12)		1,367	12.5 (23)	1,194	4.9 (69)	1,402	59.3 (42)
Lenoir	1,083	10.8 (77)	6.6 (53)		1,043	14.3 (47)	894	4.9 (69)	1,072	43.2 (78)
Lincoln	1,084	9.5 (53)	7.1 (63)		1,066	14.2 (46)	914	4.0 (42)	1,083	56.8 (47)
Macon	931	8.6 (31)	5.2 (15)		916	12.4 (20)	802	4.7 (64)	885	60.5 (37)

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one county's rate to other counties. Rank 1 = best rate.
(4) <2500 g.
(5) >4000 g.
(6) <37 weeks gestation.
(7) 37 or more weeks gestation and <2500 g.
* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 8B
2005 - 2007 Pregnancy Nutrition Surveillance ⁽¹⁾
North Carolina
Comparison of Infant Health Indicators by County
Combined 3 Years

Comparison of Infant Health Indicators ^(2, 3)									
County	Birthweight			Preterm (6)		Full Term LBW (7)		Ever Breastfed	
	Low (4)		High (5)						
	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Madison	475	7.2 (12)	6.9 (56)	469	10.9 (8)	418	2.2 (4)	472	80.1 (10)
Martin	177	11.9 (84)	9.6 (93)	175	14.3 (47)	150	5.3 (82)	120	86.7 (7)
Mcdowell	555	12.3 (87)	4.3 (9)	549	18.6 (87)	447	5.6 (87)	549	37.9 (82)
Mecklenbur	10,194	9.4 (50)	5.8 (32)	10,085	13.1 (29)	8,768	3.7 (30)	9,787	60.8 (33)
Mitchell	228	7.5 (14)	5.7 (26)	221	12.7 (26)	193	3.1 (13)	223	61.4 (31)
Montgomery	458	7.4 (13)	7.9 (77)	443	13.8 (38)	382	3.1 (13)	451	58.5 (46)
Moore	937	9.4 (50)	6.2 (46)	905	13.1 (29)	786	4.6 (60)	931	50.3 (65)
Nash	1,857	10.4 (71)	6.9 (56)	1,810	14.4 (50)	1,550	4.1 (47)	1,839	51.5 (62)
New Hanove	2,927	9.8 (58)	7.1 (63)	2,915	14.0 (42)	2,508	3.7 (30)	2,910	53.9 (57)
Northhampt	447	13.0 (93)	2.9 (1)	437	21.3 (93)	344	4.9 (69)	433	13.2 (93)
Onslow	5,208	7.7 (17)	8.5 (86)	5,155	11.9 (15)	4,542	3.1 (13)	5,086	63.4 (28)
Orange	1,169	6.8 (10)	7.9 (77)	1,137	12.4 (20)	996	3.1 (13)	1,102	71.9 (20)
Pamlico	101	5.9 (4)	10.9 (95)	97	* *	84	* *	97	* *
Pasquotank	815	7.5 (14)	4.9 (11)	809	14.5 (53)	692	3.0 (11)	773	10.3 (94)

- (1) Reporting period is January 1 through December 31.
- (2) Excludes records with unknown data or errors.
- (3) Rank compares one county's rate to other counties. Rank 1 = best rate.
- (4) <2500 g.
- (5) >4000 g.
- (6) <37 weeks gestation.
- (7) 37 or more weeks gestation and <2500 g.
- * Percentages are not calculated if <100 records are available for analysis after exclusions.

Table 8B
2005 - 2007 Pregnancy Nutrition Surveillance ⁽¹⁾
North Carolina
Comparison of Infant Health Indicators by County
Combined 3 Years

Comparison of Infant Health Indicators ^(2, 3)									
County	Birthweight			Preterm (6)		Full Term LBW (7)		Ever Breastfed	
	Low (4)		High (5)						
	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Pender	888	9.7 (56)	7.3 (67)	883	13.7 (36)	762	3.9 (38)	853	54.7 (54)
Perquimans	193	7.8 (18)	4.1 (5)	188	12.2 (17)	165	4.8 (67)	192	15.6 (92)
Person	550	10.0 (63)	9.3 (91)	540	17.8 (81)	444	5.0 (74)	545	47.2 (72)
Pitt	3,128	12.6 (90)	5.7 (26)	3,037	17.9 (82)	2,491	5.1 (76)	3,093	50.7 (64)
Polk	117	12.0 (86)	8.5 (86)	115	7.8 (2)	106	8.5 (94)	117	73.5 (18)
Randolph	2,528	7.5 (14)	8.1 (83)	2,504	13.6 (35)	2,163	2.7 (7)	2,492	68.9 (23)
Richmond	1,379	8.0 (21)	6.1 (44)	1,347	13.7 (36)	1,163	2.9 (9)	1,371	40.3 (80)
Robeson	3,285	10.3 (67)	5.6 (24)	3,216	16.7 (74)	2,678	5.2 (81)	3,263	35.3 (84)
Rockingham	1,792	9.0 (41)	5.9 (35)	1,778	14.1 (44)	1,528	4.2 (52)	1,788	48.7 (67)
Rowan	2,735	9.7 (56)	6.4 (51)	2,679	14.6 (55)	2,289	3.8 (35)	2,723	56.5 (49)
Rutherford	464	8.6 (31)	7.5 (71)	454	13.0 (28)	395	4.1 (47)	461	59.9 (40)
Sampson	1,626	8.5 (30)	6.2 (46)	1,604	14.4 (50)	1,373	3.4 (24)	1,608	35.2 (85)
Scotland	1,126	12.4 (88)	5.2 (15)	1,101	15.5 (64)	930	5.1 (76)	1,117	35.5 (83)
Stanly	951	8.6 (31)	5.3 (18)	941	12.3 (19)	825	3.9 (38)	901	77.4 (14)

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one county's rate to other counties. Rank 1 = best rate.
(4) <2500 g.
(5) >4000 g.
(6) <37 weeks gestation.
(7) 37 or more weeks gestation and <2500 g.
* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 8B
2005 - 2007 Pregnancy Nutrition Surveillance ⁽¹⁾
North Carolina
Comparison of Infant Health Indicators by County
Combined 3 Years

Comparison of Infant Health Indicators ^(2, 3)									
County	Birthweight			Preterm ⁽⁶⁾		Full Term LBW ⁽⁷⁾		Ever Breastfed	
	Low ⁽⁴⁾	High ⁽⁵⁾		Number	%	Number	%	Number	%
	Number	% (Rank)	% (Rank)						
Stokes	668	11.1 (80)	7.5 (71)	653	15.6 (65)	551	5.1 (76)	665	54.1 (56)
Surry	1,404	9.0 (41)	7.1 (63)	1,386	14.3 (47)	1,187	4.0 (42)	1,363	52.9 (59)
Swain	306	6.2 (6)	11.1 (96)	301	11.0 (9)	268	1.5 (2)	294	94.2 (2)
Transylvan	244	9.4 (50)	4.1 (5)	241	10.0 (6)	217	5.1 (76)	242	87.2 (5)
Tyrrell	112	14.3 (94)	3.6 (3)	107	18.7 (88)	87	* *	111	55.0 (53)
Union	1,762	8.2 (24)	6.9 (56)	1,742	16.3 (73)	1,458	2.5 (6)	1,657	73.8 (17)
Vance	1,159	12.8 (91)	5.3 (18)	1,132	20.1 (90)	905	6.0 (89)	973	65.8 (25)
Wake	11,351	8.1 (23)	7.7 (76)	11,155	14.6 (55)	9,529	3.3 (22)	9,657	89.4 (4)
Warren	444	14.4 (95)	5.0 (12)	432	22.0 (95)	337	7.1 (93)	438	40.4 (79)
Washington	357	10.9 (78)	7.0 (60)	353	20.4 (92)	281	4.6 (60)	355	46.2 (73)
Watauga	320	6.9 (11)	5.9 (35)	316	9.2 (3)	287	3.5 (25)	306	81.4 (9)
Wayne Coun	1,715	8.8 (37)	7.9 (77)	1,685	13.4 (32)	1,459	3.2 (18)	1,678	59.4 (41)
Wilkes	1,233	9.5 (53)	5.7 (26)	1,230	15.8 (66)	1,036	4.2 (52)	1,218	58.9 (43)
Wilson Cou	1,895	9.3 (47)	4.5 (10)	1,871	17.2 (75)	1,550	4.5 (57)	1,881	45.6 (74)

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one county's rate to other counties. Rank 1 = best rate.
(4) <2500 g.
(5) >4000 g.
(6) <37 weeks gestation.
(7) 37 or more weeks gestation and <2500 g.
* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 8B
2005 - 2007 Pregnancy Nutrition Surveillance ⁽¹⁾
North Carolina
Comparison of Infant Health Indicators by County
Combined 3 Years

Comparison of Infant Health Indicators ^(2, 3)									
County	Birthweight			Preterm (6)		Full Term LBW (7)		Ever Breastfed	
	Low (4)		High (5)						
	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Yadkin	728	8.8 (37)	5.6 (24)	723	15.8 (66)	609	3.1 (13)	726	77.5 (13)
Yancey	173	9.8 (58)	5.8 (32)	172	13.4 (32)	149	2.7 (7)	170	94.1 (3)
Unknown Counties	764	10.6 (75)	5.4 (21)	749	14.0 (42)	644	5.1 (76)	736	61.0 (32)
North Carolina	159,040	9.4	6.6	156,595	14.7	133,536	4.0	154,097	59.6

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one county's rate to other counties. Rank 1 = best rate.
(4) <2500 g.
(5) >4000 g.
(6) <37 weeks gestation.
(7) 37 or more weeks gestation and <2500 g.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

